



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: C12N 15/82, A01H 5/00, C12P 21/00, A01H 5/10, C12N 15/62, 15/56, 5/10	A2	(11) International Publication Number: WO 95/14099 (43) International Publication Date: 26 May 1995 (26.05.95)
(21) International Application Number: PCT/US94/13179 (22) International Filing Date: 15 November 1994 (15.11.94) (30) Priority Data: 08/153,563 16 November 1993 (16.11.93) US (60) Parent Application or Grant (63) Related by Continuation US 08/153,563 (CIP) Filed on 16 November 1993 (16.11.93) (71) Applicant (for all designated States except US): THE REGENTS OF THE UNIVERSITY OF CALIFORNIA [US/US]; 22nd floor, 300 Lakeside Drive, Oakland, CA 94612-3550 (US). (72) Inventor; and (75) Inventor/Applicant (for US only): RODRIGUEZ, Raymond, L. [US/US]; 3017 Anza Avenue, Davis, CA 95616 (US). (74) Agent: ROBBINS, BERLINER & CARSON; 5th floor, 201 North Figueroa Street, Los Angeles, CA 90012-2628 (US).		(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LT, LU, LV, MD, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: PROCESS FOR PROTEIN PRODUCTION IN PLANTS (57) Abstract <p>This invention provides for the secretion of heterologous protein in plant systems. In particular, this invention provides for the production of heterologous proteins by malting of monocot plant seeds. The heterologous genes are expressed during germination of the seeds and isolated from a malt. Also disclosed are chimeric genes, vectors and methods relating to the present invention. Protein production by cell culture techniques is also described.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

PROCESS FOR PROTEIN PRODUCTION IN PLANTS

FIELD OF THE INVENTION

5 The field of this invention is production of polypeptides in germinating plant seeds employing the malting process. The invention also relates to plant expression vectors, chimeric genes and plant cell culture methods.

10 BACKGROUND OF THE INVENTION

The ability to clone and produce a wide range of proteins from diverse sources became feasible with the advent of recombinant technology. The selection of expression hosts for commercial biotechnology
15 proteins is based on the economics of fermentation and purification as well as the ability of the host to accomplish the post-translational modifications needed for full biological activity of the recombinant protein. Some of these
20 post-translational modifications include: signal peptide processing, pro-peptide processing, protein folding, disulfide bond formation, glycosylation, gamma carboxylation and beta-hydroxylation. Some of the economic factors influencing the choice of an
25 expression host include: rates of biomass production, equipment costs, medium composition and expense, processes for protein recovery and purification, product yields, and the potential for contamination.

Much of the early work in biotechnology was
30 directed toward the expression of recombinant or "heterologous" proteins in prokaryotes, like *Escherichia coli* and *Bacillus subtilis*. Such work in prokaryotes provided ease of genetic manipulation, growth of the organisms in batch culture and the
35 possibility of large-scale fermentation.

E. coli can perform signal peptide processing, protein folding, and disulfide bond formation. However, it cannot secrete proteins extracellularly glycosylate, gamma carboxylate, beta hydroxylate or process propeptides. *B. subtilis* suffers from the same limitations *E. coli* except that it is capable of extracellular secretion.

Total production costs from bacteria are also high because of problems with product recovery, purification, and the inability of bacteria to perform many of the post-translational modifications mentioned above. Furthermore, *E. coli* and other bacteria are pathogens and contaminants, such as, pyrogens and endotoxins, must be removed from the recombinantly produced protein. In addition, extensive post-purification chemical and enzymatic treatments (e.g., to refold the protein into an active form) can be required to obtain biologically active protein.

Because proteins are not secreted from prokaryotes, like *E. coli*, such cells must be disrupted for product recovery. The subsequent release of bacterial contaminants and other proteins make product purification more difficult and expensive. Because purification accounts for up to 90% of the total cost of producing recombinant proteins in bacteria, proteins, like tissue Plasminogen Activator (tPA), can cost several thousand dollars per gram to produce from *E. coli*.

Because of the many inadequacies associated with prokaryotic hosts, the biotechnology industry has looked to eukaryotic hosts like mammalian cell tissue culture, yeast, fungi, insect cells, and transgenic animals, to properly and efficiently express recombinant proteins. However, these hosts can

suffer from any or all of the following disadvantages: expensive fermentation, low yields, secretion problems, inappropriate modifications in protein processing, high operating costs, difficulties in scaling up to large volumes, and/or contamination that either kills the host culture or makes product purification more expensive. For these reasons, existing eukaryotic hosts are unable to provide high-volume, low-cost protein production of recombinant proteins.

For most of those proteins requiring extensive post-translational modifications for therapeutic and/or functional activity, mammalian cell culture is the most common alternative to *E. coli*. Although mammalian cells are capable of correctly folding and glycosylating bioactive proteins, the quality and extent of glycosylation can vary with different culture conditions among the same host cells. Furthermore, mammalian culture has extremely high fermentation costs (60 - 80% of total production expense), requires expensive media, and poses safety concerns from potential contamination by viruses and other pathogens. Yields are generally low, for example, in the range of 0.5 - 1.5% of cellular protein, or up to about 300 - 400 milligrams per liter.

Yeast, fungi, insect cells and transgenic animals are currently being used as alternatives to mammalian cell culture. Yeast, however, produces incorrectly glycosylated proteins that have excessive mannose residues and generally limited eukaryotic processing. Further, although the baculovirus insect cell system can produce high levels of glycosylated proteins, these are not secreted - making purification complex and expensive. Fungi represent

the best current system for high-volume, low-cost production, but they are not capable of expressing many target proteins. Transgenic animals are subject to lengthy lead times to develop herds with stable genetics, high operating costs, and contamination by animal viruses.

The biochemical, technical and economic limitations on existing prokaryotic and eukaryotic expression systems has created substantial interest in developing new expression systems for recombinant proteins. Plants represent the most likely alternative to existing systems because of the advantageous economics of field-grown crops, the ability to synthesize proteins in storage organs like tubers, seeds, fruits and leaves and the ability of plants to perform many of the post-translational modifications previously described. However, existing plant expression systems suffer from low yield (< 1.5% of total cellular protein).

Furthermore, expression of the target protein occurs in the open field (in roots, stems, leaves, fruits and seeds), thus making it difficult to prevent the recombinant protein from entering the food and feed chain. This is an issue of much concern to government regulatory agencies.

Although the use of plant cell culture to express proteins has been discussed, the lack of knowledge about the genetics and biochemistry of plant gene expression and secretion has precluded this system from being developed into a commercially feasible one.

RELEVANT LITERATURE

The potential for the use of plant cell cultures to product proteins has been described by Zenk,

Phytochemistry 30:3861-3863 (1991). Descriptions of the rice amylase genes may be found in Huang, et al., *Proc. Natl. Sci. U.S.A.* 89:7526-7530 (1992); Huang, et al., *Plant, Molecular Biology* 14:655-668 (1990);
5 Huang, et al., *Nucleic Acids Research* 18:7007-7014 (1990); Huang, et al., *Gene* 111:223-228 (1992); Rodriguez, et al.: Organization Structure and Expression of the Rice α -Amylase Multigene Family. *Second International Rice Genetics Symposium. Rice*
10 *Genetics* 11:417-429 (1990); Sutliff, et al., *Plant Molecular Biology* 16:579-591 (1991). The promoter sequences for the rice amylase genes are described in Huang, et al., *Nucleic Acids Research* 18:7007-7014 (1990). Descriptions of plant protein signal
15 peptides may be found in addition to the references described above in Vaulcombe, et al., *Mol. Gen. Genet.* 209:33-40 (1987); Chandler, et al., *Plant Molecular Biology* 3:407-418 (1984); Rogers, J. *Biol. Chem.* 260:3731-3738 (1985); Rothstein, et al., *Gene*
20 55:353-356 (1987); Whittier, et al., *Nucleic Acids Research* 15:2515-2535 (1987); Wirsel, et al., *Molecular Microbiology* 3:3-14 (1989); Yu, et al., *Gene* 122:247-253 (1992).

A description of the regulation of plant gene
25 expression by the phytohormone, gibberellic acid and secreted enzymes induced by gibberellic acid can be found in R.L. Jones and J. MacMillin, *Gibberellins: in: Advanced Plant Physiology*, Malcolm B. Wilkins, ed., 1984 Pitman Publishing Limited, London, pp.
30 21-52. References that describe other metabolically-regulated genes: Sheen, *Plant Cell*, 2:1027-1038(1990); Maas, et al., *EMBO J.* 9:3447-3452 (1990); Benkel and Hickey, *Proc. Natl. Acad. Sci.* 84:1337-1339 (1987)

SUMMARY OF THE INVENTION

The present invention has two aspects. In the first aspect, the invention describes the production of high levels of recombinant proteins by seeds during the malting process. The invention includes a method of producing a protein or polypeptide in a monocot plant seed. Monocot plant seeds typically contain an endosperm surrounded by an aleurone or scutellar epithelium layer. Embryo development takes place within the endosperm. In the method, seeds are provided that contain a chimeric gene having at least the following components: (i) a transcriptional regulatory region inducible during seed germination, (ii) a heterologous first DNA sequence encoding the protein, and (iii) a second DNA sequence encoding signal polypeptide. The second DNA sequence is operably linked to the transcriptional regulatory region and the first DNA sequence. The signal polypeptide encoded by the second DNA sequence is in translation-frame with the protein and is effective to facilitate secretion of the protein across the aleurone or scutellar epithelium layer into the endosperm. The seeds are malted under conditions to induce expression of the transcriptional regulatory region and production of the protein or polypeptide of interest.

In this method, the leader sequence may be omitted from the chimeric gene construct if secretion of the protein of interest is not required, for example, for production of a mash or malted seed product intended for veterinary use.

In the above method, the seed may be obtained from a cereal plant, including, but not limited to wheat, rice, oats, rye, corn, sorghum, millet or

barley. Barley and rice seeds are preferred embodiments.

5 The transcriptional regulatory region of the chimeric gene used in the practice of the present invention can be obtained, for example, from one of the following groups: α -amylase genes, sucrose synthase genes and sucrose-6-phosphate synthetase genes. Examples of α -amylase genes include the following genes obtained from rice and their homologs
10 in other monocots: RAmY1A (SEQ ID NO:2), RAmY3B, RAmY3C, RAmY3D, HV18 (SEQ ID NO:1) and RAmY3E. In one embodiment of the invention the RAmY3D gene or homologs thereof is used.

15 Expression of the transcription regulatory region of the chimeric gene can be specifically regulatable by a small molecule. Examples of such small molecules include plant hormones, cytokines and metabolites. Exemplary small molecules include, but are not limited to, abscisic acid, gibberellic acids, indoleacetic acids, kinetins, butyric acid, oxalic
20 acid, acetic acid, okadacic acid or arachidonic acid. In one embodiment, the transcription regulatory region is from the RAmY3D gene, or a homolog thereof, and the small molecule is abscisic acid.

25 The malting step of the present method can also include the addition or removal of a small molecule in an amount effective to stimulate production of the protein.

Other embodiments of the transcriptional
30 regulatory region of the chimeric gene includes regulatory regions sucrose synthase genes or sucrose-6-phosphate synthetase genes. With these promoters useful regulatory small molecules include sugar (e.g., glucose or sucrose), sugar-phosphates, or
35 other sugar-derivative molecules.

The protein produced by the method of the present invention can be selected from any number of sources. Exemplary classes of proteins or polypeptides include, but are not limited to, the following: enzymes, antibodies, growth factors, cytokines, hormones, or antigens (e.g., vaccines). Specific proteins or polypeptides include, but are not limited to, the following: α -antitrypsin, antithrombin 3, fibrinogen, human serum albumin, factor VIII, granulocyte colony-stimulating factor and granulocyte macrophage colony-stimulating factor. The protein may also be an industrial protein (e.g., xylanase, oxidoreductase, peroxidase, glucanase, α -amylase, phytase or glucose oxidase).

The signal polypeptide (or signal sequence) can be obtained from a number of sources. Exemplary gene-sources for the signal sequence are α -amylase genes, sucrose synthase genes and sucrose-6-phosphate synthetase genes. Examples of α -amylase genes include the following genes obtained from rice and their homologs in other monocots: RAmY1A (SEQ ID NO:2), RAmY3B, RAmY3C, RAmY3D, HV18 (SEQ ID NO:1) and RAmY3E.

The method of the present invention can also include isolation or partial purification of the protein or polypeptide of interest.

In another embodiment, the method of the invention includes separation of the embryo from the endosperm of the seed. In this embodiment, the embryo is (i) separated from the other seed parts, and (ii) germinated under conditions that induce expression of the transcriptional regulatory region and production of the protein. The seeds used in the malting process may have one or more chimeric genes, where the genes have the same or different

transcription regulatory regions. The method may include separating the embryos and germinating them under conditions that (i) induce expression of the transcriptional regulatory region of the second
5 chimeric gene, and (ii) production of the polypeptide of the second chimeric gene. The transcriptional regulatory region can, for example, be selected from those described above. The method may also include separating embryos of the seed from the endosperm,
10 contacting the endosperm with the plant hormone which induces expression of the expression cassette, and isolating the heterologous protein from the endosperm.

Also included in the present invention are the
15 germinated seed, malted seed and mash products by the above-described method. These products may be directly used in veterinary applications where purification of the expressed protein or polypeptide is not required (e.g., growth hormone or vaccines).

20 In another aspect the invention includes the above-described chimeric gene constructs. Further, the invention includes plant expression vectors carrying the chimeric genes, plant cells bearing the chimeric gene or transformed by such vectors, and
25 transgenic plants bearing the chimeric gene. The transgenic plants and plant cells can, for example, be selected from the following group: rice, wheat, oats, rye, corn, sorghum, millet, and barley.

The present invention also includes transgenic
30 seeds produced by the transgenic plants of the present invention. Such transgenic seeds carry the chimeric gene of the present invention.

In the second aspect of the present invention, a
plant cell culture system is described for the
35 production of recombinant proteins. The cells for

the culture system are preferably derived from scutellar tissue. The cells used in this aspect of the present invention are transgenic and contain a chimeric gene. The chimeric gene contains at least
5 the following elements: (i) a transcription regulatory region inducible during seed germination, where expression mediated by the region is specifically regulatable by a small molecule, and
10 (ii) a heterologous DNA sequence that encodes the polypeptide, where the DNA sequence is operably linked to transcription regulatory region or promoter.

The chimeric genes can be constructed using the components described above, including the
15 transcription regulatory region, proteins or polypeptides for expression, and regulatory small molecules.

The invention includes plant expression vectors carrying the chimeric genes, plant cells bearing the
20 chimeric gene or transformed by such vectors, and transgenic plants bearing the chimeric gene. The transgenic plants and plant cells can, for example, be selected from the following group: rice, wheat, oats, rye, corn, sorghum, millet, and barley.

25 In one embodiment, the second aspect of the invention includes a method for modulating expression of a polypeptide in plant tissue cell culture. In this method the cells are the transgenic cells just described, typically from scutellar epithelium.
30 These cells, carrying the chimeric gene, are cultured under conditions that facilitate plant cell growth. Expression of the transcription regulatory region is modulated by addition or removal of the regulatory small molecule to the plant cell culture.

Cells used in this aspect of the invention are monocot cells from, for example, wheat, rice, oats, rye, corn, sorghum, millet or barley.

Modulation of expression can be accomplished by inhibition or induction. In one embodiment, the modulation is an inhibition of expression and the small molecule is a sugar or sugar-phosphate derivative. In another aspect, the modulation is an induction of expression and the small molecule is a plant hormone. Selected plant hormones can also be used to inhibit expression.

In the method, the culturing step can further include the addition or removal of sugar or sugar-derivatives in an amount effective to stimulate production of the heterologous protein. In another embodiment, modulation is the enhancement of expression of the polypeptide in plant tissue cell culture and the enhancement is accomplished by addition of a small molecule to the plant cell culture, where the small molecule specifically induces expression from the transcription regulatory region. An exemplary embodiment of this type of modulation is the use of the transcription regulatory region from the RAMY3D gene, or a homolog thereof, and where the small molecule is abscisic acid.

This aspect of the invention includes production and isolation or partial purification of the protein or polypeptide of interest.

The invention also includes the following method of producing a polypeptide in plant cells. The transgenic plant cells described above are cultured *in vitro* (i) under conditions that facilitate plant cell growth, and (ii) in the presence of a small molecule that inhibits expression from the transcription regulatory region. The small molecule

is present at a concentration effective to repress expression from the transcription regulatory region. The concentration of this small molecule is then reduced in the culture to a level that permits
5 expression from the transcription regulatory region and allows production of the polypeptide. Exemplary small molecules for the practice of this embodiment of the invention include sugars or sugar-phosphate derivatives. After removal of the inhibitory small
10 molecule, a second small molecule can be added to induce expression from the promoter present in the chimeric gene.

The invention also includes (i) an isolated DNA sequence essentially consisting of the nucleic acid
15 sequence of SEQ ID NO:1 or SEQ ID NO:2; an isolated DNA sequence essentially consisting of a transcriptional regulatory region obtained from SEQ ID NO:1 or SEQ ID NO:2; and (iii) one of these isolated DNA sequences operably linked to a
20 heterologous DNA sequence encoding a protein. Such isolated DNA sequences may also include a second DNA sequence encoding signal polypeptide. The second DNA sequence is operably linked to the transcriptional regulatory region and the heterologous DNA sequence.
25 Further, the encoded signal polypeptide is in translation-frame with the protein encoded by the heterologous DNA sequence.

These and other objects and features of the
30 invention will be more fully appreciated when the following detailed description of the invention is read in conjunction with the accompanying drawings.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1A is a full color photograph of the expression of the GUS gene in cell culture. Figure 1B is a full color photograph of the expression of the GUS gene in seed.

Figure 2 is a depiction of the plasmid construction for producing transgenic seed with a RAmY3D promoter.

Figure 3 shows two southern blot hybridization where the left panel (GUS) depicts DNA isolated from 3DG cells and probed with a GUS gene probe and the right panel is probed with a probe from the rice α -amylase gene (AMY).

Figure 4 provides the results of a fluorescent assay detecting GUS.

Figure 5 provides the results of a fluorescent assay detecting GUS from cells free of sucrose.

Figure 6A provides a comparison of G+C content for three different α -amylase promoters. Figure 6B is a table comparing sequence homology between α -amylase promoters.

Figure 7, Panel A depict the construct of two gene fusions used to express GUS in rice seed. Panels B and C are the results of southern blots demonstrating the stable transmission of the GUS gene to rice progeny.

Figure 8 Panels A and B are fluorometric measures of the expression of GUS using the H4 and E4 constructs in different transgenic rice seed.

Figure 9, Panels A and B are depictions of four fluorometric measures of the expression of GUS in four transgenic rice seed lines using both the H4 and E4 GUS constructs.

Figure 10, Panels A and B provide graphic depictions of four fluorometric measures of the

expression of GUS in four transgenic seed lines quantifying the induction of GUS by addition of gibberellic acid.

5 **DETAILED DESCRIPTION OF THE INVENTION**

I. DEFINITIONS

 "Cell culture" refers to cells and cell clusters both protoplast and callus tissue that are without differentiated cells or organs and are growing on a growth media.

 "In an amount effective" refers to an amount that is suitable to produce the desired effect in a measurable and reproducible amount.

 "Inducible" means a promoter that is turned on by the presence or absence of a cell hormone or metabolite. It includes both indirect and direct inducement.

 "Inducible during germination" refers to promoters which are substantially silent but not totally silent prior to germination but are turned on substantially (greater than 25%) during germination and development in the seed. Examples of promoters that are inducible during germination are presented below.

 "Homologous DNA" refers to DNA not introduced into the plant or plant host by recombinant means.

 "Heterologous DNA" refers to DNA which has been transfected into plant cells. Typically, heterologous DNA refers to DNA that is not originally derived from the transfected or transformed cells' genomic DNA (e.g., GUS gene sequences). A DNA sequence may also be heterologous to an adjacent DNA sequence, for example, a DNA sequence is heterologous to a promoter or transcription regulatory region, when the promoter or transcription regulatory region is not native,

that is, not naturally occurring adjacent the DNA sequence of interest.

5 A "transcription regulatory region" typically refers to nucleic acid sequences that influence and/or promote initiation of transcription, such as promoters. Promoters are typically considered to include regulatory regions, such as enhancer or inducer elements.

10 A "chimeric gene," in the context of the present invention, typically comprises a promoter sequence operably linked to non-homologous DNA sequences that encode a gene product (e.g., a RAmy3D promoter adjacent DNA sequences encoding human serum albumin). A chimeric gene may also contain further
15 transcription regulatory elements, such as transcription termination signals, as well as translation regulatory signals, such as, termination codons.

20 "Operably linked" refers to components of a chimeric gene or an expression cassette that function as a unit to express a heterologous protein. For example, a promoter operably linked to a heterologous DNA, which encodes a protein, promotes the production of functional mRNA corresponding to the heterologous
25 DNA.

A "product" encoded by a DNA molecule includes, for example, RNA molecules and polypeptides.

Two nucleotide sequences are considered to be "functionally homologous" if they hybridize with one
30 another under moderately stringent conditions, i.e. 0.1% SSC at room temperature. Typically, two homologous nucleotide sequences are greater than or equal to about 60% identical when optimally aligned using the ALIGN program (Dayhoff, M.O., in ATLAS OF
35 PROTEIN SEQUENCE AND STRUCTURE (1972) Vol. 5, National

Biomedical Research Foundation, pp. 101-110, and Supplement 2 to this volume, pp. 1-10.).

Two amino acid sequences are considered "homologous" if their amino acids are greater than or equal to about 60% identical when optimally aligned using the ALIGN program mentioned above.

"Removal" in the context of a metabolite includes both physical removal as by washing and the depletion of the metabolite through the absorption and metabolizing of the metabolite by the cells.

"Signal sequence suitable to permit the heterologous protein to be secreted across the aleurone or scutellular epithelium" refers to any naturally occurring signal sequence in monocots, dicots, animals or microorganisms that can permit a protein to be secreted from the cells across the stated organs of the monocot seed. Generally, a signal sequence is a sequence of amino acids that promotes the secretion of a protein from a cell.

"Small molecules" are typically less than about 1 kilodalton and are biological, organic, or even inorganic compounds (i.e., cisplatin). Examples of such small molecules include sugars, sugar-derivatives (including phosphate derivatives), and plant hormones (such as, gibberellic or abscisic acid).

"Specifically regulatable" refers to the ability of a small molecule to preferentially affect transcription from one promoter or group of promoters (e.g., the α -amylase gene family), as opposed to non-specific effects, such as, enhancement or reduction of global transcription within a cell by a small molecule.

"Substantially isolated" is used in several contexts and typically refers to the at least partial

purification of a protein or polypeptide away from unrelated or contaminating components (e.g., plant proteins removed from a sample containing human α -1-antitrypsin). Methods and procedures for the isolation or purification of proteins or polypeptides are known in the art.

II. PRODUCTION OF RECOMBINANT PROTEINS USING THE MALTING PROCESS.

The invention described herein has essentially two distinct aspects. In the first aspect, the present invention relates to the discovery of the regulated expression of recombinant proteins in malted cereal seeds. Malting is the process by which grain, typically barley or rice, is germinated under controlled conditions and in contained facilities to produce a product that can be used for human consumption, animal feed and the brewing of alcoholic beverages. The process begins by steeping barley seeds in 55°F water for 48 hours followed by a four-day germination of the grain in malting bins or drums.

During this time, the starchy portion of the seed, or endosperm, is converted to maltose and other sugars. Maltsters use water, air and, in some instances, phytohormones, like gibberellic acid, to control temperature and optimize the malting process. The malted grain is then kiln-dried at temperatures between 120°F and 130°F to terminate germination and remove moisture. At this point the malted grain can be stored or sold to the food, feed or brewing industries. Products of the malting process include mashes and formulated seed products. During the process of malting germinated seeds are also produced.

In the malting process, the rapid conversion of starch to sugar is accomplished by a tremendous burst of gene activity that results in the expression of a starch degrading enzyme called α -amylase. During the peak stage of germination, α -amylase is the major protein in the seed, constituting up to 60% of the total protein of the cells that surround the starchy endosperm. Alpha-amylase is secreted out of these cells and into the endosperm where it digests the starch into sugar.

Expression and secretion of α -amylase during germination is so abundant that it can be purified and sold as a research reagent for approximately \$0.10 per gram. The compositions and methods described herein allow low-cost, high-volume eukaryotic production of selected gene products, such as, proteins and polypeptides, based on the malting of transgenic seeds.

This aspect of the invention includes a method of producing a protein in a monocot plant seed. Monocot plant seeds contain an endosperm surrounded by an aleurone and scutellar epithelium layer. The embryo typically begins germination and development within the endosperm. Seeds are produced, for example, from transgenic plants, that contain a chimeric gene having at least the following components:

(i) a transcriptional regulatory region inducible during seed germination, several such regions are described in detail below, including promoters from α -amylase, sucrose synthase, and sucrose-6-phosphate synthetase genes;

(ii) a heterologous first DNA sequence encoding a gene product of interest, for example, a

polypeptide or protein. Exemplary DNA coding sequences are described below; and

(iii) a second DNA sequence encoding signal sequence where the second DNA sequence is operably
5 linked to the transcriptional regulatory region and the first DNA sequence. Further, the signal polypeptide sequence is (a) in translation-frame with the protein or polypeptide of interest and (b) effective to facilitate secretion of the protein or
10 polypeptide across said aleurone or scutellar epithelium layer into the endosperm.

These seeds are malted under conditions to induce expression of the transcriptional regulatory region and production of the protein or polypeptide
15 of interest.

The plants (including organs, seeds, tissues and cells) used in the process of the present invention are derived from monocots, particularly the members of the taxonomic family known as the *Gramineae*. This
20 family includes all members of the grass family of which the edible varieties are known as cereals. The cereals include a wide variety of species such as wheat (*Triticum* sps.), rice (*Oryza*, sps.) barley (*Hordeum* sps.) oats, (*Avena* sps.) rye (*Secale* sps.),
25 corn (*Zea* sps.) and millet (*Pennisetum* sps.).

Plant cells or tissues derived from the members of the *Gramineae* and are transformed with expression constructs (i.e., plasmid DNA into which the gene of interest has been inserted) using a variety of
30 standard techniques (e.g., electroporation, protoplast fusion or microparticle bombardment). In one embodiment, the expression construct includes a transcription regulatory region (promoter) whose transcription is specifically regulated by the
35 presence or absence of a small molecule.

In one embodiment, the gene encoding the recombinant protein is placed under the control of a metabolically regulated promoter. Metabolically regulated promoters are those in which mRNA synthesis or transcription, is repressed or induced by sugars or sugar derivatives. Examples of metabolically regulated promoters include those that transcribe some of the cereal α -amylase genes and sucrose synthase genes.

Another expression construct uses a hormonally regulated promoter to achieve expression of the recombinant protein in the germinated or malted seed. Hormonally regulated promoters are those in which mRNA synthesis or transcription, is repressed or induced by small molecules, like phytohormones such as gibberellic acid or abscisic acid. Other such small molecules include, but are not limited to, indoleacetic acids, kinetins, butyric acid, oxalic acid, acetic acid, okadaic acid and arachidonic acid.

Examples of hormonally regulated promoters include those that transcribe some of the cereal α -amylase genes. The promoters relevant to this application include, but are not limited to the following: those controlling the expression of the rice (*Oryza sativa*) α -amylases genes, *RAmy1A*, *RAmy3D* and *RAmy3E*; the barley α -amylase gene promoter, *HV18*; and the sucrose synthase and sucrose-6-phosphate-synthetase (SPS) promoters from rice and barley.

Expression constructs utilize additional regulatory DNA sequences (i.e., signal peptide sequences and preferred translational start codons) to promote efficient translation and extracellular secretion of the target protein. By fusing the genes for recombinant proteins to this array of regulatory

DNA sequences, the expression of recombinant proteins in germinated transgenic seeds is placed under the transcription and secretion control of a metabolically regulated or hormonally regulated promoter.

Cells or tissues or derived from cereal plants can be transformed singly or together (i.e., co-transformation) with the expression constructs. From such transformed cells transgenic plants can be regenerated. These transgenic plants are grown, allowed to produce seeds and the recombinant protein encoded by the expression construct can be recovered from malted transgenic seeds.

The principle of using different cereal α -amylase promoters to express a recombinant protein in transgenic seeds is illustrated in Figure 1B. In this figure, the gibberellic acid-induced promoter for the RAmy1A gene, was used to express the bacterial reporter gene, gusA, in rice (Example 2). The gusA gene encodes the enzyme, beta-glucuronidase (GUS), that produces a blue chromophore in tissues expressing the gene. This chromophore can be easily detected using a histochemical staining method. In transgenic rice seeds containing the RAmy1A promoter/GUS fusion, the blue chromophore increases up to six days of germination.

Using the chimeric genes, vectors and methods of the present invention, cereal species such as rice, corn, wheat, oats, rye, barley and various grasses can be genetically engineered to express a wide range of recombinant proteins. By combining the technology of the present invention with well-established production methods (e.g., crop cultivation, malting, and product recovery), recombinant protein can be efficiently and economically produced for the

biopharmaceutical, industrial processing, animal health and bioremediation industries.

5 The fact that this expression system does not require the use of genetic elements derived from animal or plant pathogens should facilitate regulatory acceptance.

A. PLANT EXPRESSION VECTORS

10 Expression vectors for use in the present invention comprise a chimeric gene (or expression cassette), designed for operation in plants, with companion sequences upstream and downstream from the expression cassette. The companion sequences will be of plasmid or viral origin and provide necessary
15 characteristics to the vector to permit the vectors to move DNA from bacteria to the desired plant host.

The basic bacterial/plant vector construct will preferably provide a broad host range prokaryote replication origin; a prokaryote selectable marker; and, for *Agrobacterium* transformations, T DNA
20 sequences for *Agrobacterium*-mediated transfer to plant chromosomes. Where the heterologous gene is not readily amenable to detection, the construct will preferably also have a selectable marker gene
25 suitable for determining if a plant cell has been transformed. A general review of suitable markers for the members of the grass family is found in Wilbanks and Dons, 1993, *Plant Mol. Biol. Reptr*, 11(2):165-185.

30 Sequences suitable for permitting integration of the heterologous sequence into the plant genome are also recommended. These might include transposon sequences, and the like, for homologous recombination, as well as Ti sequences which permit

random insertion of a heterologous expression cassette into a plant genome.

5 Suitable prokaryote selectable markers include resistance toward antibiotics such as ampicillin or tetracycline. Other DNA sequences encoding additional functions may also be present in the vector, as is known in the art.

10 The constructs of the subject invention will include the expression cassette for expression of the protein(s) of interest. Usually, there will be only one expression cassette, although two or more are feasible. The recombinant expression cassette contains, in addition to the heterologous protein encoding sequence, at least the following elements:
15 a promoter region, plant 5' untranslated sequences, initiation codon depending upon whether or not the structural gene comes equipped with one, and transcription and translation termination sequences. Unique restriction enzyme sites at the 5' and 3' ends
20 of the cassette allow for easy insertion into a pre-existing vector. These elements are discussed in detail below.

1. HETEROLOGOUS CODING SEQUENCES.

25 The heterologous coding sequence may be for any protein of interest, either prokaryotic or eukaryotic, particularly eukaryotic. The gene providing the desired product will particularly be those genes associated with large volume products.
30 Therefore, products of particular interest include but are not limited to enzymes, such as chymosin, proteases, polymerases, saccharidases, dehydrogenases, nucleases, glucanase, glucose oxidase, α -amylase, oxidoreductases (such as fungal peroxidases and laccases), xylanases, phytase,
35

cellulase, hemicellulase, and lipase. More specifically, the invention can be used to produce enzymes such as those used in detergents, rennin, horse radish peroxidase, amylases from other plants, soil remediation enzymes, and other such industrial proteins.

Other proteins of interest are mammalian proteins. Such proteins include, but are not limited to blood proteins (such as, serum albumin, Factor VII, Factor VIII (or modified Factor VIII), Factor IX, Factor X, tissue plasminogen factor, Protein C, von Willebrand factor, antithrombin III, and erythropoietin), colony stimulating factors (such as, granulocyte colony-stimulating factor (G-CSF), macrophage colony-stimulating factor (M-CSF), and granulocyte macrophage colony-stimulating factor (GM-CSF)), cytokines (such as, interleukins), integrins, addressins, selectins, homing receptors, surface membrane proteins (such as, surface membrane protein receptors), T cell receptor units, immunoglobulins, soluble major histocompatibility complex antigens, structural proteins (such as, collagen, fibroin, elastin, tubulin, actin, and myosin), growth factor receptors, growth factors, growth hormone, cell cycle proteins, vaccines, fibrinogen, thrombin, cytokines, hyaluronic acid and antibodies.

While for the most part, the product will be a peptidic product, genes may be introduced which may serve to modify non-peptidic products produced by the cells. These proteins, fragments thereof, usually of at least about 30 amino acids, fused combinations, mutants, and synthetic proteins, whether the proteins may be synthetic in whole or in part, so far as their

sequence in relation to a natural protein, may be produced.

The present invention also provides the advantage that polypeptide useful for veterinary use, such as, vaccines and growth hormones, may be produced by the malting process of the present invention. The products of the malting reaction, containing the polypeptide of interest, can then be formulated into mash product or formulated seed product directly useful in veterinary applications.

2. SIGNAL SEQUENCES

Also included in chimeric genes used in the practice of the method of the present invention are signal secretion sequences. In addition to encoding the protein of interest, the chimeric gene also encodes a signal peptide that allows processing and translocation of the protein, as appropriate. The chimeric gene typically lacks any sequence that might result in the binding of the desired protein to a membrane.

Typically, the transcriptional regulatory region (*i.e.*, the transcription initiating region) is derived from a gene whose product is expressed and translocated during germination. By employing the signal peptide homologous to such a transcriptional regulatory region, translocation of the protein of interest is achieved. In this way, the protein(s) of interest are translocated from the cells in which they are expressed and may be efficiently harvested.

While it is not required that the protein be secreted from the cells in which the protein is produced, this facilitates the isolation and purification of the recombinant protein. Table 1 provides a list of known signal sequences from wheat,

barley and rice. Typically, these signal sequences facilitate secretion of proteins expressed in seeds across the aleurone or scutellar epithelium layer into the endosperm of the seed.

5

Table 1

Signal peptides of α -amylase genes

B- , barley α -amylase genes. W- , wheat α -amylase genes; #, signal peptide cleavage site determined by protein sequencing; *, predicted signal peptide cleavage site; /, intron splice site; ".", space inserted to maximize the alignment.

	<u>Genes</u>	<u>References</u>
15	RAmy1A	Plant Molecular Biology. 14:655-668 (1990).
	RAmy1B	Nucleic Acids Research. 18:7007-7014 (1990).
	aAmy10-c	Gene. 122:247-253 (1992).
20	W-Amy1/13 Mol.	Gen. Genet. 209:33-40 (1987).
	W-2128	Gene. 55:353-356 (1987).
	B-pM/C	Plant Molecular Biology. 3:407-418 (1984) and Journal of Biological Chemistry. 260:3731-3738 (1985).
25	B-gKAmy141	Plant Molecular Biology. 9:3-17 (1987).
	RAmy2A	Gene. 111:223-228 (1992).
	W-Amy2/54	Mol. Gen. Genet. 209:33-40 (1987).
	B-clone E	Plant Molecular Biology. 3:407-418 (1984) and J. of Biological Chemistry 260:3731-3738 (1985).
30	B-gKAmy155	Plant Molecular Biology. 3:407-418 (1984) and Plant Molecular Biology. 9:3-17 (1987).

	B-Amy32b	Plant Molecular Biology. 3:407-418 (1984) and Nucleic Acids Research 15:2515-2535 (1987).
5	RAmy3A	Plant Molecular Biology. 16:579-591 (1991).
	RAmy3B	Plant Molecular Biology. 16:579-591 (1991).
	RAmy3C	Plant Molecular Biology. 16:579-591 (1991).
10	RAmy3D	Nucleic Acids Research. 18:7007-7014 (1990).
	RAmy3E	Nucleic Acids Research. 18:7007-7014 (1990).
	W-Amy3/33	Mol. Gen. Genet. 209:33-40 (1987).
15	Taka-amylase	Molecular Microbiology. 3:3-14 (1989).

Those of skill can routinely identify new signal peptides. Plant signal peptides typically have a tripartite structure, with positively-charged amino acids at the N-terminal end, followed by a hydrophobic region and then the cleavage site within a region of reduced hydrophobicity. The conservation of this mechanism is demonstrated by the fact that cereal α -amylase signal peptides are recognized and cleaved in foreign hosts such as *E. coli* and *S. cerevisiae*.

The flexibility of this mechanism is reflected in the wide range of polypeptide sequences that can serve as signal peptides. Thus, the ability of a sequence to function as a signal peptide may not be evident from casual inspection of the amino acid sequence. Methods designed to predict signal peptide cleavage sites identify the correct site for only about 75% of the sequences analyzed. (See Heijne Gv:

Cleavage-site motifs in protein targeting sequences. In: J. K. Setlow (eds) *Genetic Engineering*, Vol. 14. Plenum Press, New York (1992)).

5 Although, sequence homology is not always present in the signal peptides, hydrophilicity plots demonstrate that the signal peptides of these genes are relatively hydrophobic.

3. PROMOTERS

10 (a) EXEMPLARY TRANSCRIPTION REGULATORY REGIONS

 The preferred transcription regulatory or promoter region is chosen so as to be relatively silent, except during seed germination. For example, the expression level in the seed cells is at least
15 about 20 times the expression level in other plant tissue during the growth of the plant. This type of transcriptional regulation can be achieved in various ways, including the following: (i) by using the
20 5'-non-coding region associated with a protein which is produced solely or substantially solely during seed germination, or (ii) by using the regulatory portion of such a transcriptional initiation region in conjunction with a different RNA polymerase binding region.

25 In referring to a "substantial absence of expression" at times other than seed germination, it is intended that expression be very low or non-existent except during seed germination. That is, the expression of the protein of interest,
30 encoded by the chimeric gene, is low so as (i) to not affect the growth of the plant or expend significant plant resources, (ii) to not diminish the vigor of the plant growth, and (iii) to allow for the plant and mash to be used for its intended purpose,
35 depending on the protein of interest.

A number of proteins are normally secreted across either the aleurone or scutellum during seed germination and seed elongation. Some examples of secreted plant enzymes induced by gibberellic acid include, but are not limited to the following: α -amylase, protease, ribonuclease, β -glucanase, esterase, acid phosphatases (such as p-nitrophenyl, phosphatase, ATPase, phytase, naphthol AS-B1 phosphatase, and GTPase), pentosanase, endoxylanase, xylopyranosidase, arabinofuranosidase, glucosidase, and peroxidase.

In view of the teachings of the present specification, one of skill in the art can recognize and implement useful promoter/signal sequence combinations for the practice of the present invention. Because many of the useful sequences are evolutionarily related, the conserved sequences facilitate the identification of new promoters and signal sequences useful in the practice of this invention.

Standard nucleic acid hybridization technology can be used to probe libraries of other monocots, using previously identified promoters and signal sequences, to identify "homologs" in these monocots. For example, a gene having a promoter region of interest is selected, a probe specifically hybridizable to the gene is chosen, and traditional cross-hybridization experiments are performed under varying solution stringencies. This method has been employed using the rice RAmY1A as a probe to identify a homolog promoter in barley; the barley promoter for HV18 (SEQ ID NO:1).

Polymerase chain reaction technology (PCR; Mullis, K.B., U. S. Patent No. 4,683,202, issued 28 July 1987; Mullis, K.B., et al., U. S. Patent No.

4,683,195, issued 28 July 1987) can also be used to identify homologs of known promoters, for example, amplify unknown promoter using PCR primers able to bind to conserved regions of a selected promoter or signal sequence. Examples of conserved sequences in the rice amylase promoter regions are provided in Table 2. The sequences for the rice promoters were reported in Huang N., et al., 1990, *Nucl. Acids Res.* 18:7007-7014 (1990) and the taka promoter from *Aspergillus oryzae* was reported in Wirsal, 1989, *Molecular Microbiology*, 3:3-14.

Table 2
Conserved sequences in the *RAmy3D*, *RAmy3E* and
Taka-amylase promoters

31 bp <i>RAmy3D</i>	GAGACCGGGCCCCGACGCGGCGGACGCGGCG	SEQ ID NO:3
	++++ + ++ +++ + ++ ++ ++++++	
31 bp <i>RAmy3E</i>	GAGAGCTCGCGCCGCTCGATCGGCGCGGCG	SEQ ID NO:4
11 bp <i>RAmy3D</i>	TTCCGGCTTGC	SEQ ID NO:5
	++ ++++++	
11 bp <i>RAmy3E</i>	TTGCGGCTTGC	SEQ ID NO:6
Taka-amylase	CGGCCCGTCCGC	SEQ ID NO:7

The "+" symbols indicate positions at which the *RAmy3D* and *RAmy3E* sequences are identical.

The situation in rice is demonstrative. In rice, the α -amylase isozymes are encoded by a family of nine genes (Table 3). They are referred to as *RAmy1A*, *1B*, *1C*, *2A*, *3A*, *3B*, *3C*, *3D* and *3E*. The Rice α -Amylase genes are classified into three subfamilies (*RAmy1*, *RAmy2*, and *RAmy3*) (See Huang, et al., 1992, *Proc. Natl. Acad. Sci. USA.* 89:7526-7530 and Huang N, et al., 1990, *Plant Molecular Biology*, 14:655-668) based on DNA sequence similarities to α -amylase gene subfamilies in other cereal species. Eight members of the α -amylase gene family in rice have been

isolated and characterized. A partial cDNA sequence, presumably corresponding to *RAmy1C*, has been reported in Yu S-M, et al., 1992, *Gene*. 122:247-253. The α -amylase genes have been mapped to five different chromosomes in rice.

Table 3
Alpha-amylase gene expression in rice tissues

5	Gene Names	Germinated Seedlings	Root	Leaf	Developing Seeds	Cultured Cells
	All Genes	100 ^a	1	3	4	65
	RAmy1A	++++ ^b	++++	++++	+++	++
	RAmy1B	-	-	-	-	+
	RAmy1C	- ^c	NA ^d	NA	NA	NA
10	RAmy2A	+	+	+	+	+
	RAmy3A	+	-	-	-	++
	RAmy3B	++	NA	NA	NA	NA
	RAmy3C	++	NA	NA	NA	NA
	RAmy3D	++	++	++	-	++++
15	RAmy3E	+++	+++	+++	++++	+++

^a Relative mRNA levels for all α -amylase genes are normalized to the level of expression observed in germinated seedlings by Northern blot hybridization.

^b Relative levels of mRNA for each gene as estimated from Northern blot hybridization or RNA-PCR experiments. The amount of PCR product is indicated from the highest (++++) to the lowest (+). Minus signs (-) indicate that no product was observed in the RNA-PCR reaction.

^c Lack of expression based on restriction digest of RNA-PCR products.

^d NA = Not Available

^e The nine α -amylase genes represent all members of the gene family found in rice cv. IR36.

Embodiments of the present invention include promoter/signal sequence combinations of RAmy 1A, 3D and 3E, where expression is at a high level during germination. The 5'-non-coding region of 3E is

characterized by a region conserved with 3D, which is a GC-rich sequence of 31 bases and contains two CGGC repeats. There is also an 11 base sequence which is conserved which contains a single copy of the CGGC sequence.

5

The 3D and 3E α -amylase genes are subject to suppression of expression by sugars, particularly sucrose, glucose, fructose, and maltose. Thus, during cell fermentation, premature expression of the desired product can be avoided by employing a sugar, particularly, sucrose, in the growth medium. Thus, sucrose may be used as a carbon source by the cells and, when the sucrose is exhausted, expression of the desired proteins will be initiated.

10

15

Other transcription initiation regulatory regions that may be employed include those that are induced by sugar such as sucrose synthase.

20

Complex transcriptional initiation regions can be employed by using the regulatory portion of one transcriptional initiation region with the RNA polymerase binding region of a different gene. In this way, expression can be regulated while providing for a high level of expression.

25

A preferred embodiment of the present invention uses a promoter that is regulated during germination. For example, the hormones abscisic acid (ABA) and gibberellic acid (GA) play important regulatory roles in control of α -amylase gene expression in cereal seeds. ABA, which is synthesized during grain filling, acts as a negative regulator of transcription for α -amylase and many other genes. ABA levels drop in the mature desiccated grain, thus relieving the inhibition of α -amylase and other genes required for germination.

30

Obviously up-regulation is desired for over expression of heterologous proteins and GA mediated promoters are a desired embodiment. The prevailing model for GA regulation during cereal seedling
5 development involves the diffusion of GA from the embryo to the aleurone layer. GA then induces the synthesis of hydrolytic enzymes such as α -amylase. In rice, GA stimulates α -amylase gene expression in aleurone tissues.

10 But not all α -amylase promoters are induced by GA and not all are inducible when present in undifferentiated cells, such as those used in culture or when removed from the intact seed. For example *RAmy3D* and *RAmy3E* in rice callus and cultured cells
15 are unaffected by GA. No significant change in levels of *RAmy3D* and *RAmy3E* expression was detected in rice callus treated with GA and there are reports that callus treated with paclobutrazol, an inhibitor of gibberellin biosynthesis, produced the same amount
20 of α -amylase protein as untreated callus.

Finally, callus cultures derived from seeds of the GA-deficient dwarf mutant, cv. Tan-ginbozu, produced the same levels of α -amylase gene expression with or without exogenous GA treatment. This
25 suggests that *Ramy3D* and *Ramy3E* gene expression in the scutellum and in cultured cells is independent of GA regulation.

The GA independent promoters appear to be missing a short sequence that is present in GA
30 inducible promoters. DNA sequence comparisons have identified four short, conserved sequences in the cereal α -amylase promoters. The TATA Box (CTATAAATAG) is the binding site required for RNA polymerase II to initiate transcription. The Pyrimidine Box (YCTTTTY) and Box I (TATCCAT) may be involved in the
35

developmental regulation of the genes in the scutellum and aleurone.

The GARE Box (GA-Responsive Element) (TAACRRA) is required for GA-induction and ABA-repression of α -amylase gene expression. The GARE Box (GA-Responsive Element) in the *RAmy1A* gene (genomic clone 10Sg2) is located at base -143 relative to the transcription start site. Expression of the *RAmy1A* gene (cDNA clone pOS103) is stimulated 50-100 fold by exogenous GA. The absence of GARE Box sequences in the promoters of the rice *RAmy3D* and *RAmy3E* genes is consistent with the GA-independent expression of these genes as discussed above.

Alternatively the agent inducing expression can be a metabolite which is either a sugar or phosphorylated sugar. For example, *RAmy3D* gene expression is metabolically regulated in rice embryo tissues. Evidence for this is based on studies in which seeds were moistened to initiate seedling development and harvested after 0 to 48 hours of incubation. Embryos dissected out of these seeds had low levels of expression for *RAmy3D*. This pattern of expression was reversed if embryos were first removed from the seed at time zero and incubated in water for 0 to 48 hours. Under these conditions, *RAmy3D* expression increased to five times the level observed in the intact rice seed.

Addition of sugar to the incubation medium used for the isolated embryos restored normal expression of the *RAmy3D* gene (see examples). A number of sugars, including sucrose, glucose, fructose and maltose, were able to repress *RAmy3D* gene expression in isolated embryos (Karrer and Rodriguez, 1992, *The Plant Journal*, 2:517-523).

In rice cell suspension cultures, α -amylase enzyme activity increases after the depletion of sucrose from the medium. This increase is consistent with the pattern of α -amylase mRNA accumulation, which also increases dramatically after the culture medium is depleted of sugar. Cells transferred to sugar-free medium begin to produce elevated levels of total α -amylase mRNA within four hours. (See Yu S-M, et al., 1991, *J Biol. Chem.* 266:21131-21137 (1991).

Dot blot hybridization and gene-specific probes have been used to demonstrate that sugar controls the expression of both *RAmy3D* and *RAmy3E* in cultured cells. Cells were subcultured into medium with 1%, 3%, 6% or 12% sucrose. RNA isolated from cells cultured for five days showed that *RAmy3D* and *RAmy3E* expression was repressed at the higher sugar concentrations. Expression of both of these genes was induced in the cells cultured in 1% sucrose, presumably after sugar was depleted from the medium by cell growth. Expression of the *RAmy1A* and *RAmy3A* genes was unaffected by these treatments. Thus, expression of the *RAmy3D* and *RAmy3E* genes is metabolically regulated by the concentration of sugar in the culture medium.

To confirm that this regulation is acting at the level of transcription, the *RAmy3D* promoter was linked to the GUS reporter gene. The amount of GUS enzyme activity produced by the cells provides a convenient measure of the expression of the engineered *RAmy3D* promoter. The *RAmy3D* promoter/GUS gene construct was co-electroporated into protoplasts with another plasmid containing the gene for hygromycin-resistance. Southern blot hybridization showed that hygromycin-resistant cell lines contained

the *RAmy3D*/GUS gene construct. GUS activity was reduced in transformed cell lines grown at elevated sucrose concentrations (see examples). GUS activity increased starting eight hours after the cells were transferred to sugar-free medium. These data demonstrate that the GUS reporter gene is being regulated by the *RAmy3D* promoter in cell culture just as the endogenous *RAmy3D* gene is regulated in the rice seed.

The promoters of the *RAmy3D* and *RAmy3E* genes have little sequence similarity, but there are two conserved sequences in their promoters that may be involved in the metabolic regulation of these genes. A GC-rich sequence of 31 bases in *RAmy3D* (Table 2) contains two CGGC repeats (underlined). This tandem repeat structure within the 31 base sequence is similar to the DNA binding sites for Sp1, a mammalian transcription regulatory protein.

An eleven-base sequence (Table 2), which is conserved in both the *RAmy3D* and the *RAmy3E* promoter, contains a single copy of the CGGC sequence. A tandem duplication of the CGGC sequence is also found in the promoters of the Taka-amylase genes of *Aspergillus oryzae*. These CGGC sequences are found in the 87 base region of the Taka-amylase promoters (from position -377 to -290) that has been implicated in the metabolic regulation of these genes.

(b) GENERAL SOURCES OF PROMOTER AND SIGNAL SEQUENCES

In view of the guidance of the present disclosure and by using a number of standard procedures, one of skill can identify suitable promoters and signal sequences in addition to those described above for use in this invention. While the gene can be amplified directly from a mRNA extract

using PCR, the first step is generally to produce a genomic or cDNA library.

In brief, genomic or cDNA libraries are prepared according to standard techniques as described, for instance, in Sambrook, et al., *Molecular Cloning - A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York 1989. To construct genomic libraries, large segments of genomic DNA are generated by random fragmentation and are ligated with vector DNA to form concatemers that can be packaged into the appropriate vector. Two kinds of vectors are commonly used for this purpose, bacteriophage lambda vectors and cosmids. Alternatively, genomic libraries can be purchased from commercial sources (e.g., Clontech, Palo Alto, CA).

In the present invention, cDNA libraries enriched, for example, for α -amylase secreting mRNA sequences are used to screen for the desired genes. Preparation of appropriately enriched cDNA would involve the use of plant organs over expressing and secreting α -amylase, e.g., aleurone layers. Other organs would include roots, stems, leaves and panicles. Briefly, mRNA from select tissue is isolated and cDNA is prepared. Short chains of oligo d-T nucleotides are hybridized with the poly-A tails of the mRNA and serve as a primer for the enzyme, reverse transcriptase, which synthesizes a complementary DNA (cDNA) strand.

The cDNA can be optionally enriched for the desired sequences using subtraction hybridization procedures by labelling the cDNA and hybridizing it with mRNA from tissue that does not express the desired mRNA according to the procedures. *Proc. Natl. Acad. Sci. U.S.A.* 81:2194-2198 (1984).

Unreacted cDNA is isolated and used to prepare a library for screening. To do this, a second DNA strand is synthesized using the first cDNA strand as a template. Linkers are added to the double-stranded cDNA for insertion into a plasmid or λ phage vector for propagation in *E. coli*.

Identification of clones harboring the desired sequences is performed by either nucleic acid hybridization or immunological detection of the encoded protein, if an expression vector is used. Typically, oligonucleotide probes specific for the gene of interest are used.

After identification of a cDNA corresponding to the gene of interest, the cDNA molecule can be used as a probe of a genomic DNA, containing the gene of interest. Corresponding genomic clones are identified. Genomic clones isolated with this approach will contain gene sequences that include the promoter, coding region, introns and terminators.

Alternatively, the probes specific to the gene of interest can be used to directly probe genomic DNA libraries from a selected plant. Sequences homologous to the probe can be isolated by standard techniques, including hybridization screening or polymerase chain reaction.

Oligonucleotide and cDNA probes useful for identification of other promoters and signal sequences can also be prepared from conserved regions of related genes in other species. By comparing the nucleotide sequences of the known proteins, one simply identifies conserved sequences in the genes and uses those sequences as probes or as PCR primers to locate homologous sequences in genomic or cDNA libraries of other plants. A number of references compare regions of nucleotide homology and amino acid

identity regarding secreting genes and they are provided below. Such conserved sequences can be used to isolate other genes having a hormonal or other metabolite responsive promoter.

5 Probes, typically used to identify related but heretofore unknown target sequences, can be hybridized under stringent conditions to ensure that the sequences are in fact related. Typically, stringent conditions suitable for finding related
10 sequences would be performing the hybridization at a melting temperature (T_m) of between -15°C to -20°C .

(c) CLONING OF THE DESIRED DNA SEQUENCES

Once the DNA encoding the desired sequences has
15 been located, sufficient quantity of the gene must be generated to facilitate subsequent recombinant manipulations. Although the sequences can be directly amplified by PCR, they are most commonly replicated in an intermediate bacterial host. Most
20 commonly in a bacteria of the genera *Escherichia*, *Bacillus* and *Streptomyces*. Cloning for amplification of intermediate vectors is most preferred in *E. coli* because that organism is easy to culture and more fully understood than other species of prokaryotes.

25 Sambrook, *supra*, contains methodology sufficient to conduct clonings in *E. coli*. Strain HB101 is a useful strain which is typically grown on Luria broth (LB) with glucose, Difco's Antibiotic Medium #2 and M9 medium supplemented with glucose and acid-hydrolyzed casein amino acids. Strains with resistance to
30 antibiotics are maintained at the drug concentrations described in Sambrook, *supra*.

Transformations are performed according to the method described by Morrison, D.A. (1977),
35 *J. Bacteriol.*, 132:349-351; or by Clark-Curtiss,

J.E. and Curtiss, R., 1983, in *Methods in Enzymology*, 101:347-362, Wu, R., Grossman, L. and Moldave, K., eds., Academic Press, New York. Representative vectors include pBR322 and the pUC series which are
5 available from commercial sources.

4. TRANSCRIPTION AND TRANSLATION TERMINATORS

The expression cassettes or chimeric genes of the present invention typically have a
10 transcriptional termination region at the opposite end from the transcription initiation regulatory region. The transcriptional termination region may normally be associated with the transcriptional initiation region or from a different gene. The
15 transcriptional termination region may be selected, particularly for stability of the mRNA to enhance expression. Illustrative transcriptional termination regions include the NOS terminator from *Agrobacterium* Ti plasmid and the rice α -amylase terminator.

20 Polyadenylation tails, Alber and Kawasaki, 1982, *Mol. and Appl. Genet.* 1:419-434 are also commonly added to the expression cassette to optimize high levels of transcription and proper transcription termination, respectively. Polyadenylation sequences
25 include but are not limited to the *Agrobacterium* octopine synthetase signal, Gielen, et al., *EMBO J.* 3:835-846, 1984 or the nopaline synthase of the same species Depicker, et al., *Mol. Appl. Genet.* 1:561-573, 1982.

30 Since the ultimate expression of the desired gene product will be in a eukaryotic cell (e.g., a member of the grass family), it is desirable to determine whether any portion of the cloned gene contains sequences which will be processed out as
35 introns by the host's splicosome machinery. If so,

site-directed mutagenesis of the "intron" region may be conducted to prevent losing a portion of the genetic message as a false intron code, Reed and Maniatis, *Cell* 41:95-105, 1985.

5

5. TRANSFORMATION OF PLANT CELLS

(a) DIRECT TRANSFORMATION

Vectors containing a chimeric gene of the present invention can introduced into plant cells by a variety of techniques. These vectors may include selectable markers for use in plant cells (such as, the nptII kanamycin resistance gene). The vectors may also include sequences that allow their selection and propagation in a secondary host, such as, sequences containing an origin of replication and a selectable marker. Typical secondary hosts include bacteria and yeast. In one embodiment, the secondary host is *Escherichia coli*, the origin of replication is a *colE1*-type, and the selectable marker is a gene encoding ampicillin resistance. Such sequences are well known in the art and are commercially available as well (e.g., Clontech, Palo Alto, CA; Stratagene, La Jolla, CA).

The vectors of the present invention may also be modified to intermediate plant transformation plasmids that contain a region of homology to an *Agrobacterium tumefaciens* vector, a T-DNA border region from *Agrobacterium tumefaciens*, and chimeric genes or expression cassettes (described above). Further, the vectors of the invention may comprise a disarmed plant tumor inducing plasmid of *Agrobacterium tumefaciens*.

Vectors useful in the practice of the present invention may be microinjected directly into plant cells by use of micropipettes to mechanically

transfer the recombinant DNA. Crossway, *Mol. Gen. Genet*, 202:179-185, 1985. The genetic material may also be transferred into the plant c/ell by using polyethylene glycol, Krens, et al., *Nature*, 296, 72-74, 1982.

Another method of introduction of nucleic acid segments is high velocity ballistic penetration by small particles with the nucleic acid either within the matrix of small beads or particles, or on the surface, Klein, et al., *Nature*, 327, 70-73, 1987 and Knudsen and Muller, 1991, *Planta*, 185:330-336 (teaching particle bombardment of barley endosperm to create transgenic barley).

Yet another method of introduction would be fusion of protoplasts with other entities, either minicells, cells, lysosomes or other fusible lipid-surfaced bodies, Fraley, et al., *Proc. Natl. Acad. Sci. USA*, 79, 1859-1863, 1982.

The vector may also be introduced into the plant cells by electroporation. (Fromm, et al., *Proc. Natl Acad. Sci. USA* 82:5824, 1985). In this technique, plant protoplasts are electroporated in the presence of plasmids containing the gene construct. Electrical impulses of high field strength reversibly permeabilize biomembranes allowing the introduction of the plasmids. Electroporated plant protoplasts reform the cell wall, divide, and form plant callus.

(b) VECTORED TRANSFORMATION

A common vector method of introducing the vector into plant cells is to infect a plant cell with *Agrobacterium tumefaciens* previously transformed with the gene. Under appropriate conditions known in the

art, the transformed plant cells are grown to form shoots or roots, and develop further into plants.

5 *Agrobacterium* a representative genus of the gram-negative family *Rhizobiaceae*. Its species are responsible for plant tumors such as crown gall and hairy root disease. In the dedifferentiated tissue characteristic of the tumors, amino acid derivatives known as opines are produced and catabolized. The bacterial genes responsible for expression of opines are a convenient source of control elements for
10 chimeric expression cassettes.

Heterologous genetic sequences, such as the chimeric genes of the present invention, can be introduced into appropriate plant cells, by means of
15 the Ti plasmid of *Agrobacterium tumefaciens*. The Ti plasmid is transmitted to plant cells on infection by *Agrobacterium tumefaciens*, and is stably integrated into the plant genome. J. Schell, *Science* 237: 1176-1183, 1987.

20 Ti plasmids contain two regions essential for the production of transformed cells. One of these, named transferred DNA (T-DNA), is transferred to plant nuclei and induces tumor formation. The other, termed virulence region, is essential for the
25 transfer of this T-DNA but is not itself transferred. The transferred DNA region, which transfers to the plant genome, can be increased in size by the insertion of the gene encoding group 3 LEA proteins without its ability to be transferred being affected.

30 A modified Ti plasmid, in which the tumor-causing genes have been deleted, can be used as a vector for the transfer of the gene constructs of this invention into an appropriate plant cell.

35 Construction of recombinant Ti plasmids in general follows methods typically used with the more

common bacterial vectors such as pBR322. Additional use can be made of accessory genetic elements sometimes found with the native plasmids and sometimes constructed from foreign sequences. These may include but are not limited to "shuttle vectors",
5 Ruvkun and Ausubel, 1981, *Nature* 298:85-88, promoters, Lawton, et al., 1987, *Plant Mol. Biol.* 9:315-324 and structural genes for antibiotic resistance as a selection factor, Fraley, et al.,
10 *Proc. Natl. Acad. Sci.* 80:4803-4807, 1983.

Species which are a natural plant host for *Agrobacterium* may be transformable *in vitro*. Monocotyledonous plants, and in particular, cereals and grasses, are not natural hosts to *Agrobacterium*.
15 Attempts to transform them using *Agrobacterium* have been unsuccessful until recently. Hooykas-Van Slogteren, et al., *Nature* 311:763-764, 1984. There is growing evidence now that certain monocots can be transformed by *Agrobacterium*. Using novel
20 experimental approaches that have now become available, cereal and grass species may now be transformed.

Promoters directing expression of selectable markers used for plant transformation (e.g., nptII) should operate effectively in plant hosts. One such
25 promoter is the nos promoter from native Ti plasmids, Herrera-Estrella, et al., *Nature* 303:209-213, 1983. Others include the 35S and 19S promoters of cauliflower mosaic virus, Odell, et al., *Nature*
30 313:810-812, 1985, and the 2' promoter, Velten, et al., *EMBO J.* 3, 2723-2730, 1984.

6. PLANT REGENERATION

After determination of the presence and
35 expression of the desired gene products, whole plant

regeneration is desired. Plant regeneration from cultured protoplasts is described in Evans, et al., *Handbook of Plant Cell Cultures*, Vol. 1: (MacMillan Publishing Co. New York, 1983); and Vasil I.R. (ed.), *Cell Culture and Somatic Cell Genetics of Plants*, Acad. Press, Orlando, Vol. I, 1984, and Vol. III, 1986.

All plants from which protoplasts can be isolated and cultured to give whole regenerated plants can be transformed by the present invention so that whole plants are recovered which contain the transferred gene. It is known that practically all plants can be regenerated from cultured cells or tissues, including but not limited to all major species of sugarcane, sugar beet, cotton, fruit and other trees, legumes and vegetables, and monocots.

Some suitable plants include, for example, species of wheat, rice, oats, rye, corn, sorghum, millet and barley.

Means for regeneration vary from species to species of plants, but generally a suspension of transformed protoplasts containing copies of the heterologous gene is first provided. Callus tissue is formed and shoots may be induced from callus and subsequently rooted.

Alternatively, embryo formation can be induced from the protoplast suspension. These embryos germinate as natural embryos to form plants. The culture media will generally contain various amino acids and hormones, such as auxin and cytokinins. It is also advantageous to add glutamic acid and proline to the medium, especially for such species as corn and alfalfa. Shoots and roots normally develop simultaneously. Efficient regeneration will depend on the medium, on the genotype, and on the history of

the culture. If these three variables are controlled, then regeneration is fully reproducible and repeatable.

5 The mature plants, grown from the transformed plant cells, are selfed and non-segregating, homozygous transgenic plants are identified. The inbred plant produces seed containing the chimeric gene of the present invention. These seeds can be malted to produce the protein or polypeptide of
10 interest.

The inbreds according to this invention can be used to develop hybrids or novel varieties embodying the desired traits. Such plants would be developed using traditional selection type breeding.

15 The transgenic seeds produced by the method of the present invention can be formed into seed products.

B. ANTISENSE APPLICATIONS

20 In addition to the above indicated genes, one may also have constructs which provide for inactivation of endogenously expressed genes. Of particular interest is the inactivation of genes that are expressed during germination and seedling
25 elongation. These genes may include one or more of the amylases, e.g., RAmy3B, RAmy3C, RAmy3E or homologs thereof.

Inactivation of gene expression may be achieved in a number of ways. The most convenient is the use
30 of an anti-sense sequence, where the anti-sense sequence may be complementary to any portion of the mRNA, including both the non-coding and coding regions. Normally, the anti-sense sequence will be at least about 30 nt, more usually at least about 50
35 nt, and may be up to or greater than the sequence of

the mRNA to which the anti-sense sequence is complementary.

5 In one embodiment, the 3'-terminal sequence of the anti-sense sequence is selected to provide for mRNA stability, there being a number of sequences which are known to destabilize the mRNA which can be avoided.

10 The transcription initiation region for the anti-sense sequence may be constitutive or inducible. A relatively strong promoter may be employed, such as, the 35S CMV promotor, the RUBSICO promoter, or beta-conglycinin promoter. Preferably, the transcription initiation region is inducible so as to be induced during the malting process. To enhance
15 the transcription of the anti-sense sequence, one may use various enhancers associated with other promoters to increase the rate of transcription of the anti-sense sequence.

20 It is not necessary that all expression of one or more proteins naturally produced during malting is inhibited, it being sufficient that there be at least about a 10%, preferably at least about a 25% reduction in expression, so as to increase the proportion of the desired protein in the malting
25 product. Enhancers which find use include the 35S CMV enhancer, and the introns of the alcohol dehydrogenase gene of maize.

III. THE MALTING PROCESS

30 The malting process is a multi-step process. The first step is steeping. During steeping seed is immersed in or sprayed with water to increase the moisture content of the seed to between 35-45%. This initiates germination. Steeping typically takes
35 place in a steep tank which is typically fitted with

a conical end to allow the seed to flow freely out. The addition of compressed air to oxygenate the steeping process is an option. The temperature is controlled at approximately 22°C depending on the seed.

After steeping, the seed is transferred to germination compartments. The seed is either wet or dry transferred. The germination bin contains air saturated with water and is under controlled temperature and air flows. The typical temperatures are between 12-25°C and germination is permitted to continue for from 3 to 7 days.

Where the heterologous protein is operably linked to a inducible promoter requiring a metabolite such as sugar or plant hormone, this metabolite is added, removed or depleted from the steeping water medium and/or is added to the water saturated air used during germination. The seed absorbs the aqueous medium and begins to germinate expressing the heterologous protein. The medium may then be withdrawn and the malting begun, by maintaining the seeds in a moist temperature controlled aerated environment. In this way, the seeds may begin growth prior to expression, so that the expressed product is less likely to be partially degraded or denatured during the process. Other components included in the imbibition medium may be plant hormones, such as gibberellic acid, generally in an amount from about 2.5 to 100 μM .

Where the promoter is induced by sugar, glucose or sucrose can be added to the imbibition media during steeping or during germination. The sugar concentration may range up to about 12 weight percent of the medium.

More specifically, the temperature during the imbibition or steeping phase will be maintained in the range of about 15-25°C, while the temperature during the germination will usually be about 20°C.

5 The time for the imbibition will usually be from about 2 to 4 days, while the germination time will usually be an additional 2 to 10 days, more usually 3 to 7 days. Usually, the time for the malting does not exceed about ten days. The period for the

10 malting can be reduced by using plant hormones during the imbibition, particularly gibberellic acid.

Germinated seed produced by this method may be used to produce germinated seed products.

To achieve maximum production of recombinant

15 protein from malting, the malting procedure will be modified to accommodate de-hulled and de-embryonated seeds. The hulls and embryos are dehulled and de-embryonated using standard means which include rollers, other mechanical means of breaking the

20 intact embryos free of hull and endosperm. Screening is typically used to separate the embryos from unwanted seed debris. Isolated transgenic embryos are germinated in steeping water containing CaCl_2 (approx 10 mM).

25 In the absence of sugars from the endosperm, there is expected to be a 5 to 10 fold increase in RAmy3D promoter activity and thus expression of the heterologous protein. Alternatively when embryoless-half seeds are incubated in 10mM CaCl_2 and 5 μM

30 gibberellic acid, there is a 50 fold increase in RAmy1A promoter activity.

In this system, recombinant proteins under the control of RAmy1A (HV18 or other homologs) and RAmy3D promoters are secreted into the medium. Specialized

35 malting bins or steep tanks may be used. The embryos

and embryoless-half seeds are mechanically disrupted to release any secreted protein between cells and tissues. The mixture is suspended in a buffer solution to retrieve soluble proteins. Conventional
5 protein isolation and purification methods are then used to purify the recombinant protein. Parameters of time, temperature pH, oxygen, and volumes are adjusted through routine methods to optimize expression and recovery of heterologous protein.

10 An optional step is kilning the germinated seed. Kilning is a low temperature drying procedure that reduces moisture concentration to 4-6%. Temperatures during kilning are between 40-85°C. Typically the
15 lower temperatures of less than 60°C are used until the seed have a moisture content of between about 10 and 20%. Final drying is at higher temperatures of above 60°C.

The kiln-dried material (or malted seeds) may then be used directly, for example, where the
20 recombinant protein is useful in veterinary applications (e.g., animal feed containing a an increased protein value, growth hormone or vaccine). The mash may be formulated into a mash product.

The mash may also be processed by mechanical
25 disruption of the seeds and bringing the total protein into solution. The cellular debris may then be separated by any convenient means, such as settling, centrifugation, filtration, or the like. The supernatant or filtrate will normally include
30 from 1 to 40 weight percent of the desired product of total protein in the medium, preferably at least about 30 weight percent.

Where the desired product is not water soluble, one may need to extract the desired product with a
35 convenient solvent or use another process which

allows for solubilization and/or extraction of the product without loss of the desired activity of the product or which allows for renaturation.

5 After isolation of the protein of interest from the aqueous medium, one may then purify the product in accordance with conventional ways. Since the product will be a substantial portion of the total protein present in the mixture, frequently being present in the greatest percentage of any individual
10 protein, purification is greatly simplified. Furthermore, contaminants in the product after purification are not likely to be of physiological concern for many of the applications of the products, including therapeutic applications.

15 By providing for malting, seeds can be germinated under conditions where a desired product can be produced in the germinated seeds to provide for a high proportion of the total protein in the malting mash being the desired protein. By breaking
20 the cells, separating the cellular debris from the protein, and isolating the supernatant from the mash, the protein may be easily isolated and purified, being a major component of the total protein in the medium.

25 As distinct from other methods of producing proteins, the subject method provides for high levels of economic production of proteins in a crude form which can be easily purified. The system lacks the potential for the production of endo and exotoxins,
30 which is of concern with prokaryotes. The system allows for storage under ambient conditions without significant loss of seed viability or product loss. The product can be produced on demand. In this way, proteins can be produced in accordance with need,

where the source of the protein can be conveniently and safely stored.

5 In one embodiment, the malted transgenic seeds of the present invention, containing a protein of interest, can be used as a malted seed feed product for animals. The malted transgenic seeds can be mixed with normal (i.e., non-transgenic) seeds to obtain the desired concentration of the protein of interest in an animal feed product.

10 The following is an example of protein production by the malting process of the present invention. Varieties of rice containing the human gene for α -1-antitrypsin (i.e., transgenic rice) are inbred to produce homozygous lines. These transgenic
15 lines are cultivated in the field and mature seeds are harvested using conventional agricultural practices.

The transgenic seeds are transported to a malting facility (malthouse) where they are soaked
20 (steeped) in water for 48 hours. The seeds are then transferred to malting bins where they are allowed to germinate from 2 to 4 days. During germination, the temperature and humidity of the seed bed is monitored to ensure ideal germination conditions. In some
25 cases, chemicals such as the plant hormone, gibberellic acid, may be added to enhance the expression of the α -1-antitrypsin gene which is under the control of a gibberellic acid inducible promoter (e.g., RAmy1A, RAmy3C, HV18).

30 After optimum germination and expression of the α -1-antitrypsin gene have been achieved, the seeds are mashed (for example, by gentle grinding) to disrupt tissues and remove the hulls from the seeds. The seed mash is suspended in a protein extraction
35 buffer. Such a buffer typically contains protease

inhibitors, reducing agents and a buffering agent (such as, "TRIS" or sodium or potassium phosphate).

The mash is agitated or stirred to ensure that all secreted protein is freed from tissues and cells.

5 Large particulate matter, such as hull, plant tissues, and other debris are then removed by filtration or centrifugation. The supernatant is collected and chilled to reduce proteolysis of α -1-antitrypsin.

10 The supernatant is subjected to various purification schemes used in the wet-milling industry (e.g., hydrocloning and ion exclusion chromatography) to remove un-wanted proteins and to concentrate α -1-antitrypsin. Alternatively, ammonium sulfate
15 precipitation can also be used to concentrate the α -1-antitrypsin.

Affinity- and ion-exchange chromatography can be used to purify the α -1-antitrypsin away from other proteins in the supernatant. The presence of α -1-
20 antitrypsin in the various chromatographic fractions can be detected using standard photometric assays.

In another embodiment, after the transgenic seeds are transported to a malting facility (malthouse) they are dehulled and deembryonated
25 (i.e., mechanical separation of the embryos and endosperm portions of the seed). The embryos and endosperms are separately soaked (steeped) in water for 48 hours. The seeds are treated as described above. The separated embryos are treated as follows.
30 Expression of the RAmY3D promoters is induced in the absence of sugar and/or by the addition of chemicals, such as a plant hormone, e.g., abscisic acid.

After optimum germination and expression of the α -1-antitrypsin gene have been achieved, the embryo
35 and endosperm portions are mixed and then mashed

(i.e., gentle grinding) to disrupt seed tissues. The mash is then treated as above for purification of the α -1-antitrypsin polypeptide.

5 IV. Production of Recombinant Proteins Using a Cell Culture Process.

 In the second aspect, the invention relates to the regulated expression of recombinant proteins in cereal cell culture. In one embodiment of the invention, the cells are derived from scutellar epithelium of cereal plants. The chimeric genes, vectors and methods described above may be implemented in the practice of this aspect of the invention. In this aspect, the invention includes modulating expression of a polypeptide in monocot plant tissue cell culture. Transgenic plant cells are produced that contain a chimeric gene having at least the following components:

 (i) a transcription regulatory region inducible during seed germination, where expression mediated by said region is specifically regulatable by a small molecule. Several examples of such regions and small molecule regulators are described below; and

 (ii) a heterologous DNA sequence that encodes the polypeptide, where said DNA sequence is operably linked to said promoter. The chimeric gene may also contain a signal sequence to facilitate secretion from the cell of the polypeptide encoded by the heterologous DNA sequence.

 The transgenic cells are cultured under conditions that facilitate plant cell growth. Expression of the polypeptide of interest is modulated by addition or removal of at least one small molecule to the plant cell culture.

For example, the principle of using different cereal α -amylase promoters to express a recombinant protein in tissue culture cells is illustrated in Figure 1A. In this figure, the sugar-repressible promoter for the rice α -amylase gene, RAmy3D, was
5 used to express the bacterial reporter gene, *gusA*, in rice. The *gusA* gene encodes the enzyme, beta-glucuronidase (GUS), that produces a blue chromophore in tissues expressing the gene. This
10 chromophore can be easily detected using a histochemical staining method. As can be seen in this figure, the product of *gusA* is repressed in rice cells when the culture medium contains 3% sugar.

Cells or tissues derived from cereal plants can
15 be transformed singly or together (i.e., co-transformation) with the expression constructs described above. Once integrated into the plant genome, the recombinant protein can be recovered and purified from the medium of cultured transgenic cells.

20 The vectors of this invention can be used to facilitate the expression and/or secretion of heterologous protein in cell culture. The plant cells are placed and maintained into suspension culture and induced through the variety of inducers
25 described above to produce high levels of the desired heterologous protein. The protein is then isolated using conventional technology.

Because the purifications are dramatically varied for individual proteins, it is sufficient to
30 indicate that the initial purification process will typically follow the purification process of the native protein from its host. Because the growth media of the plant suspension culture, as used in the present invention, is typically more simple than the
35 normal host environment of the protein of interest,

the purification procedures may be appropriately modified and simplified by those of skill in the art.

It is evident from the above results, that plant cells can be engineered and the cells used to propagate plants. The plant cells can be modified to provide for expression constructs that allow controlled expression of the coding sequence in the construct to provide the expression product as the major product.

By combining the technology of the present invention with well-established production methods (e.g., plant cell fermentation, crop cultivation, and product recovery), recombinant protein can be efficiently and economically produced for the biopharmaceutical, industrial processing, animal health and bioremediation industries.

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be readily apparent to those of ordinary skill in the art in light of the teachings of this invention that certain changes and modifications may be made thereto without departing from the spirit or scope of the appended claims.

The following examples are provided by way of illustration only and not by way of limitation. Those of skill will readily recognize a variety of noncritical parameters which could be changed or modified to yield essentially similar results.

EXAMPLES

GENERAL METHODS

Generally, the nomenclature and laboratory procedures with respect to standard recombinant DNA

technology can be found in Sambrook, et al.,
Molecular Cloning - A Laboratory Manual, Cold Spring
Harbor Laboratory, Cold Spring Harbor, New York 1989
and in S.B. Gelvin and R.A. Schilperoot, *Plant*
5 *Molecular Biology*, 1988. Other general references
are provided throughout this document. The
procedures therein are known in the art and are
provided for the convenience of the reader.

10

EXAMPLE 1

Expression of β -glucuronidase
under the control of the α -amylase promoter
in rice cell culture

15

A. *Initiation of scutellar callus and*
suspension cultures.

Rice seeds (*Oryza sativa* L. cv. M202) were
provided by Dr. M. Brandon (California Rice
Experimental Station). Seeds were dehulled, washed
20 three times with water, rinsed in 70% ethanol for 20
sec and then surface-sterilized in 1% sodium
hypochlorite with a few drops of Tween 20 under
vacuum for 20 min. Sterilized seeds were washed
three times with sterile distilled water. Seven
25 seeds were placed in 15 cm petri dishes containing LS
medium with 2 mg/l 2,4-D and 30 g/l sucrose. The
seeds were incubated in the dark at 28°C and checked
periodically to monitor the growth of scutellar-
derived callus. Callus formation from scutellum
30 tissue and/or embryo was visible after 5 days. After
30 to 40 days, clumps of friable calli, about 1 cm in
diameter, were saved and the remaining tissue was
discarded.

35

To initiate a suspension culture, friable calli
were gently agitated in a petri dish with liquid AA
medium as described by Thompson JA, Abdullah R and

Cocking EC, Protoplast culture of rice (*Oryza sativa* L.) using media solidified with agarose (*Plant Sci.* 47:123-133 (1986) to reduce the calli to small clusters of cells. Cell clusters from about 20-30 clumps of calli were then transferred to a 125 ml Erlenmeyer flask and the liquid was replaced with 25 ml of fresh AA medium. The flasks were incubated in the dark on a rotary platform shaker at 110 rpm and 28°C. The primary culture was sub-cultured every 4 to 5 days with repeated screening for small cell clusters. This was accomplished by passing the culture sequentially through nylon filters of 1000 μ m and 500 μ m pore size. After two months of subculture, a finely divided and rapidly growing suspension culture was obtained. This culture was subsequently maintained by weekly subculture in AA medium containing 3% sucrose.

B. Construction of the RAmy3D/GUS gene fusion

The RAmy3D promoter/GUS gene fusion shown in Figure 2 was constructed in three steps. First, a 1.5 kb *Sall* fragment containing the promoter and part of the coding region from rice genomic clone λ OSg1A as described by Huang, et al., 1990a, *Nucleic Acid Res.*, 18:7007-7014 was subcloned into pBluescript KS- to produce the plasmid pIAS1.5. The *AluI* fragment from pIAS1.5 containing 876 bp of promoter and 66 bp of 5' untranslated region was subcloned into the *EcoRV* site of "pBLUESCRIPT KS+" to form p1Alu.

Second, a plasmid containing a promoterless GUS cassette was constructed by subcloning the *HindIII/EcoRI* GUS cassette from pBl101 (Jefferson RA, 1987, Assaying chimeric genes in plants: the GUS gene fusion system. *Plant Mol. Biol. Reporter*, 5:387-405) into pUC19 to form pBl201. A pUC19

polylinker in front of the GUS coding region provides convenient cloning sites for inserting promoter fragments. Third, the *RAmy3D* promoter fragment was inserted into the promoterless GUS plasmid to produce the plasmid p3DG. The *Xba*I/*Alu*I (in *Hind*III site) promoter fragment from p1Alu was ligated into *Xba*I/*Sma*I digested pBl201. The final 11 bp of *RAmy3D* 5' untranslated region was substituted by 21 bp from the polylinker resulting in the 5.83 kb plasmid, p3DG.

The junction between the *RAmy3D* promoter and the 5' end of the GUS gene was confirmed by DNA sequencing. DNA restriction digest, DNA gel electrophoresis, ligation, transformation, plasmid DNA isolation and DNA sequencing followed standard procedures (Sambrook, et al., *Molecular Cloning - A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York 1989).

C. Protoplast isolation and DNA transformation

Three days after subculture, a 60 ml of rice suspension culture was given a ten minute, 45°C heat-treatment and transferred to two glass crystallizing-dishes. The AA medium was removed and the cells in each dish were mixed with the 20 ml of enzyme mixture (1% Cellulose RS, 1% Macerozyme R10 in CPW medium described by Thompson, et al., 1986, *Plant Sci.* 47:123-133. The cell walls were digested at 28°C for 15 hours while shaking at 45 rpm.

After digestion, the protoplasts were screened through 150, 50 and 20 μ m nylon filters, washed three times by centrifuging for 10 min at 80g, and gently resuspended in 40 ml of CPW medium. Then the suspension was adjusted to 5 million protoplasts/ml.

Two and one-half million protoplasts in 0.5 ml volume were mixed with 5 μ g of p3D2 DNA, 25 μ g of calf thymus carrier DNA and 5 μ g of pGL2 plasmid DNA carrying the CaMV 35S promoter/hph gene fusion encoding hygromycin-resistance as described by Shimamoto K, Terada R, Izawa T and Fujimoto H (Fertile transgenic rice plants regenerated from transformed protoplasts *Nature* 338:274-276 (1989)), transferred to a cuvette, placed on ice for ten min, then electroporated with a "GENE PULSER" electroporator (Bio-Rad) (300 volt/cm, 560 mF and 600 Ohms).

After being kept on ice for an additional 10 min, the protoplasts were mixed with 0.5 ml of 4X KPR medium (Thompson, et al., *Supra* 1986) and 1 ml of melted 2.4% "SEA PLAQUE" low gelling agarose, plated in Petri dishes and incubated at 28°C in the dark.

Ten days after plating, the agarose in each Petri dish was separated into four pieces and transferred into a 6 cm Petri dishes containing 5 ml of liquid KPR medium. Four days later hygromycin was added to each dish to a final concentration of 50 mg/ml. Hygromycin-resistant colonies were picked and grown in liquid AA medium to form a number of cell lines. A sample from each cell line was then assayed for GUS activity by staining the cells with X-glu (Jefferson, et al., *supra* 1987). Cell lines having GUS activity were retained. Uniform cell lines were obtained by two additional rounds of isolating cell single cell clusters coupled with selection for GUS expression. Integration of p3DG DNA into the rice genome was verified by Southern blot analysis.

D. RNA isolation and dot blot hybridization

Total RNA was isolated from cell suspension culture using a modification of the phenol/SDS procedure Ausubel FM, Brent R, Kingston RE, Moore DD, Seidman JG, Smith JA and Struhl K: Current Protocols in Molecular Biology. (1989). Approximately one gram of cultured rice cells were ground into a fine powder with sand and 5 ml of liquid nitrogen until all the liquid nitrogen evaporated. Then 2.5 ml of TLE buffer (0.2 M Tris pH. 8.2, 0.1 M LiCl, 5 mM EDTA, 20 mM sodium metabisulfite) was added and grinding continued until the sample was completely liquefied.

At this point, 0.5 ml of 10% SDS, 2.5 ml of phenol and 2.5 ml of chloroform were added to the mortar sequentially and mixed well by grinding. The sample was centrifuged at 4000 g for 15 min and the aqueous phase removed and extracted with chloroform. Total RNA was precipitated by the addition of 1/3 volume of 8 M LiCl and the mixture was allowed to sit overnight at 4°C. The RNA was harvested by centrifugation at 16,000 g for 30 min at 4°C and the RNA pellet was dissolved in 0.5 ml of double distilled water treated with diethylpyrocarbonate. The RNA was extracted once more with chloroform and then precipitated with ethanol. The RNA yield was approximately 500 µg from each gram (fresh weight) of cells.

The pre-hybridization and hybridization of α-amylase probe to the membrane under α-amylase group-specific conditions was as previously described by Huang N, Koizumi N, Reinl S and Rodriguez RL (Structural organization and differential expression of rice α-amylase genes *Nucleic Acids Res.* 18:7007-7014 (1990a) and Huang N, Sutliff TD, Litts JC and Rodriguez RL: Classification and

characterization of rice α -amylase multigene family. *Plant Mol. Biol.* 14:655-668 (1990b)).

Four different rice α -amylase genes were used as probes. The *RAmy1A* probe, a 1.6 kb *Xba*I fragment
5 from pOS103 described in O'Neill SD, Kumagai MH, Majumdar A, Huang N, Sutliff TD and Rodriguez RL (The α -amylase genes in *Oryza sativa*: characterization of cDNA clones and mRNA expression during seed
10 germination. *Mol. Gen. Genet.* 221:235-244 (1990)) cross hybridized with the closely related genes *RAmy1B* and *RAmy3C* under the stringency conditions used.

The *RAmy3D* probe, a 1.6 kb *Xba*I fragment from pOS137 [O'Neill, *supra*] was used under gene-specific
15 conditions. The *RAmy3A* probe, a 3.5 kb *Eco*RI fragment from λ OS7D (Sutliff TD, Huang N, Litts JC and Rodriguez RL: Characterization of an α -amylase multigene cluster in rice. *Plant Molecular Biology.* 16:579-591 (1991)) was used under highly stringent,
20 gene-specific conditions. The *RAmy3E* probe, a 2 kb *Hind*III fragment including the two introns exons II and III, and the 3' end as described by Huang N, Koizumi N, Reinl S and Rodriguez RL (Structural organization and differential expression of rice α -
25 amylase genes. *Nucleic Acids Res.* 18:7007-7014 (1990a)).

E. DNA isolation and Southern blot hybridization

30 Total genomic DNA was isolated using a small scale CTAB procedure described by Rogers SO and Bendich AJ (Extraction of DNA from milligram amounts of fresh, herbarium and mummified plant tissues. *Plant Mol. Biol.* 5:69-76 (1985)). Southern blot
35 analysis of transformed and un-transformed cultured

cells was the same as previously described in Huang N, Sutliff TD, Litts JC and Rodriguez RL (Classification and characterization of rice α -amylase multigene family. Plant Mol. Biol. 14:655-668 (1990b)).

The DNA probes used were the *Hind*III/*Eco*RI fragment of pBI201 (Jefferson, *Supra*) for the GUS gene and pOS103 as described in O'Neill SD, Kumagai MH, Majumdar A, Huang N, Sutliff TD and Rodriguez RL (The α -amylase genes in *Oryza sativa*: characterization of cDNA clones and mRNA expression during seed germination. Mol. Gen. Genet. 221:235-244 (1990) for rice α -amylase gene *RAmyLA*.)

F. Isolation of total protein and β -glucuronidase (GUS) activity assay

Total water soluble protein was isolated from suspension culture cells based on the procedure described by Jefferson, *supra*. A 200 mg sample of suspension culture cells was ground in a mortar and pestle for one min in the presence of sand and 0.5 ml of GUS extraction buffer (Jefferson, *supra*). The slurry was transferred to a 1.5 ml microfuge tube and the cell debris removed after 5 min of centrifugation at room temperature. The supernatant was saved as a crude extract of water soluble protein. The GUS activity was measured by the fluorometric assay procedure (Jefferson *supra*) The background level of GUS activity in control, untransformed cells was negligible.

GUS activity was also assayed in whole cultured cells by calorimetric methods (Jefferson, *supra*). Fresh or frozen cultured cells were put in either a 1.5 ml tube, a 3.5 cm Petri dish or a microtiter plate. Five volumes of sterile X-glu solution were

added to the cells. The reaction was incubated at 37°C for 30 min. or longer.

5 G. *Secretion of GUS into the media using the
 RAMy1A promoter in transgenic rice cells.*

The suspension cells are removed by filtration and the filtered media is assayed for GUS activity. The assay methods are as described above.

10 H. RESULTS

 1. *Metabolic regulation of RAMy3D and RAMy3E in
 untransformed rice suspension cultures.*

 The above experiments allow for the determination of which rice α -amylase gene(s) is metabolically regulated in rice cell cultures. Total RNA was isolated from a cell culture sampled over a period of eight days. RNA dot blots were hybridized sequentially with four different α -amylase gene probes. The mRNA levels for the RAMy1A, RAMy1B and RAMy1C genes were low and did not change significantly during the eight day period. The level of RAMy3A mRNA was also low and showed little change during the culture cycle. The levels of RAMy3D and RAMy3E mRNA were low initially, but increased significantly after five days, reaching their peak levels at 8 and 6.5 days respectively. These results are consistent with previous studies which demonstrated that RAMy3D (Group 2) and RAMy3E (Group 5) mRNA were abundantly expressed in rice cell culture while the expression of RAMy1A/RAMy1B/RAMy1C (Group 1), RAMy2A (Group 4) and RAMy3A/RAMy3B/RAMy3C (Group 3) was either low or undetectable. Other workers have found moderate expression of RAMy1A and another gene in the Ramy1 subfamily, but only in 14 day old cell cultures.

The concentration of sugar in the rice cell suspension culture medium correlates with the amount of α -amylase enzyme produced by the cells. To investigate this effect at the gene level, suspension
5 cultures, normally maintained in medium containing 3% sucrose, were subcultured into media with 1%, 3%, 6% or 12% sucrose. RNA was isolated from cells harvested at one day and at five days after subculture. RNA dot blots were hybridized with α -
10 amylase gene probes. All cells harvested after one day had approximately the same levels of α -amylase gene expression, presumably because none of the cultures had yet depleted the sucrose from the medium.

15 The mRNA levels for the *RAmy3D* and *RAmy3E* genes increased significantly after five days in the culture with 1% sucrose medium, showing induction of gene expression after the sugar was depleted from the medium. Cultures with higher initial sucrose
20 concentrations still had only low levels of *RAmy3D* and *RAmy3E* gene expression after five days. The levels of *RAmy1A*, *RAmy1B*, *RAmy1C* and *RAmy3A* gene expression changed little in any of the cultures.

Sucrose concentrations in the culture medium
25 were altered to test the effect on α -amylase gene repression. After four days of culture in medium with an initial sucrose concentration of 3%, cultures were subdivided and the sucrose concentration was increased to 6% and 12%. One of the cultures was
30 washed and resuspended in sucrose-free medium (0%). Cultures were incubated for two more days and RNA was isolated for analysis by slot blot hybridization.

RAmy1A, *RAmy1B*, *RAmy1C* and *RAmy3A* expression levels were consistently low in all subcultures.
35 *RAmy3D* gene expression was significantly reduced in

subcultures supplemented with 6% or 12% sucrose, relative to that of the sucrose-free subculture. *RAmy3E* gene expression remained high in all treatments. Thus, within two days after the addition of sucrose to the culture media, *RAmy3D* was highly repressed while *RAmy3E* expression was relatively unchanged. It is not clear to what extent these results are due to differential transcriptional control and/or differential mRNA stability.

2. Transformation of *RAmy3D* promoter/*GUS* into rice cell lines.

A *RAmy3D* promoter/*GUS* gene fusion was constructed and used to transform rice protoplasts. The plasmid p3DG contains 876 bp of *RAmy3D* 5' flanking region plus 66 bp of the 5' untranslated leader sequence linked to the *GUS* coding region (Figure 2). Plasmid p3DG was introduced into rice protoplasts by co-electroporation with the plasmid pGL2 which carries the hygromycin-resistance gene. Protoplast-derived colonies were selected on hygromycin-containing medium and tested for co-transformation with the *RAmy3D/GUS* construct by staining a few cells from each colony for *GUS* activity. Two cycles of hygromycin-resistance selection and *GUS* activity screening were used to isolate the 3DG cell line.

DNA was isolated from the 3DG cell line and from a non-transformed control cell line, digested with *Bam*HI and subjected to Southern blot hybridization (Figure 3). When the blot was probed with the *GUS* gene, a strong hybridization signal to DNA from the 3DG cell line (lanes 2-5) was observed. No hybridization was seen with DNA isolated from the control cell line (panel *GUS*, lane 1 of Figure 3).

The negative result in the GUS panel (lane 1) was not due to the lack of DNA transferred to the membrane. Equivalent amounts of DNA were detected in all lanes when the same membrane was stripped of the GUS probe and rehybridized with a probe from the rice α -amylase gene *RAmy1A* (Amy panel). These bands hybridizing to the α -amylase probe have the molecular weights predicted from the DNA sequence of the *RAmy1A* gene.

Using the GUS probe, the 3DG cell line had the same hybridization pattern before (GUS panel, lane 2) and after (lane 4) the two cycles of single cell clump selection, indicating that the plasmid DNA was stably inherited as the cells proliferated.

Two types of structural evidence indicate that the *RAmy3D/GUS* DNA is integrated into the chromosomes of the 3DG cell line. First, Southern blot analysis revealed that the GUS gene probe hybridized exclusively to undigested genomic DNA larger than the size of the p3DG plasmid (GUS panel, lanes 3 & 5).

Second, digestion of the DNA from the 3DG cell line with endonuclease *Bam*HI resulted in multiple hybridization bands (GUS panel, lanes 2 & 4). *Bam*HI does not cut within the *RAmy3D/GUS* gene construct, so each band size represents a different sized junction fragment between the unique *Bam*HI site in the p3DG plasmid and a *Bam*HI site in the adjacent chromosomal DNA.

Thus, multiple copies of the *RAmy3D/GUS* gene construct (and at least one copy of the hygromycin-resistance gene) have been integrated into the genome of the 3DG cell line. The low molecular weight bands and the faint bands of hybridization on the Southern blot probably represent fragments of the *RAmy3D/GUS* gene construct inserted into the rice genome.

3. Metabolic regulation of *RAmy3D/GUS* in transgenic rice cell lines.

Gene expression and enzyme activity for GUS was assayed in the 3DG cell line to determine whether the promoter fragment in the *RAmy3D/GUS* construct contains all of the *cis*-elements necessary for proper- expression and metabolic regulation of the gene. Dot-blot hybridization using a GUS gene probe indicated that the mRNA level from the GUS gene in 3DG cells increased as sugar was depleted from the culture medium.

The GUS enzyme assay was used to test for the expression of *RAmy3D/GUS* in response to various concentrations of sucrose in the culture medium. The 3DG cell line was subcultured into modified AA medium containing 0%, 3% or 12% sucrose. Three days later, water soluble protein was extracted and assayed for GUS by the fluorescence assay (Figure 4). The GUS activity in cells cultured with no sucrose was 65-fold higher than that of cells grown in 3% sucrose and 130-fold higher than that of cells grown in 12% sucrose. Thus, the transcriptional activity of the *RAmy3D* promoter was greatly repressed in the presence of high levels of sucrose while being highly induced under conditions of sugar deprivation.

The timing of *RAmy3D* promoter induction in response to sugar deprivation was studied by incubating 3DG cells in sucrose-free medium. There was little or no increase in GUS activity during the first eight hours of incubation (Figure 5). GUS activity increased rapidly between eight to thirty-two hours after subculturing. The expression and metabolic regulation of the *RAmy3D/GUS* gene construct resembles that of the endogenous *RAmy3D* gene.

These results are similar to those of others who observed an increase in total α -amylase mRNA beginning 4 hours after the start of sugar deprivation. Thus, the *cis*-element(s) responsible for metabolic regulation must be contained in the 942 bp promoter region on the *RAmy3D/GUS* construct.

The expression of the GUS gene product was visualized using histochemical staining methods (Jefferson, *supra*) as seen in Figure 1A where cell cultures incubated in the absence of sugar show blue staining with relatively high blue staining evident and where 3% sugar repressing the expression of the GUS gene product with relatively less blue staining is evident.

4. Promoter sequence analysis

Promoter sequences for *RAmy3D* and *RAmy3E* were compared to gain additional insight into the metabolic regulation of these genes. Two regions of sequence similarity were previously identified in the promoters of these genes. One of these regions consists of a 31 bp GC-rich sequence that is 71% identical between the *RAmy3D* and *RAmy3E* genes. This sequence is not found in the *RAmy1A* promoter (Figure 6A) or in any other rice α -amylase promoter. This sequence is found at position -264 in *RAmy3D* (Figure 6B) and contains three nearly perfect repeats of a hexanucleotide sequence composed solely of G and C residues.

The *RAmy3E* promoter contains one complete and one partial copy of the hexanucleotide repeat sequence. The tandem duplication of GC-rich hexanucleotides in the 31 bp GC-rich sequences is reminiscent of binding sites for the mammalian transcription factor Spl.

An 11 bp sequence containing part of the GC-rich hexanucleotide is also found in the *RAmy3D* and *RAmy3E* promoters (Figure 6B). These sequences may represent cis-acting elements involved in the metabolic regulation of the rice α -amylase genes. GC-rich promoter sequences have also been identified in the metabolically regulated α -amylase genes of *Aspergillus oryzae*.

10

EXAMPLE 2

Secretion of heterologous protein
across the aleurone layer of an intact rice seed
using the *RAmy1A* promoter

A. Plasmids

15

Plasmids were constructed using standard recombinant DNA methods (Ausubel, et al., 1989, *Current Protocols in Molecular Biology*, NY John Wiley and Sons and Sambrook, et al., supra, 1989). The *RAmy1A* gene of rice was chosen because of its responsiveness to GA (O'Neil, et al., 1990 *Mol. Gen Genet.*, 221, 235-224) and because it is the most active of the α -amylase genes expressed during seed germination (Karrer, et al., 1991, *Plant Mol. Biol.*, 16, 797-805).

20

Two regions of the *RAmy1A* promoter were fused to the *gus A* reporter gene to produce plasmids pH4/GUS (-748 to +31) and pE4/GUS (-232 to +31). Both promoter regions contain three conserved sequences (⁻²¹⁴CCCTTT⁻²⁰⁹, ⁻¹⁴⁷TAACAAA⁻¹⁴¹, and ⁻¹³⁰TATCCAT⁻¹²⁴) found in all the GA-responsive cereal α -amylase genes examined to date (Huang, et al., 1990). An additional pyrimidine box is present in pH4/GUS at position -312.

25

The promoter for the *RAmy1A* gene was subcloned as a 2.3 kb DNA fragment from the rice genomic DNA clone (10Sg2) into "pBLUESCRIPT M13+KS." The nucleotide sequence of this promoter has been described in Huang,

30

35

et al., 1990 Nuc. Acids Res. 18:7007-7014. The principal features of these constructs consists of the b-glucuronidase gene (*gusA*), together with the transcriptional terminator of the nopaline synthase gene from pBI101 as reported in Jefferson, 1987 *EMBO Journal* 6:3901-3907

The expression cassette was inserted into the *Sma*I site of "pBLUESCRIPT" and designated pBSGUS. *RAmy1A*(promoter)/*gusA* gene fusions were constructed by inserting restriction fragments containing the *RAmy1A* promoter into pBSGUS. The restriction fragments used to make constructs were the *Pst*I-*Hind*III fragment (-748 to +31, *pH4/GUS*) and the *Pst*I-*Eco*RI fragment (-232 to +31, *pE4/GUS*). The coordinates used to describe these restriction fragments are based on transcription start point for *RAmy1A* (Huang, et al., supra 1990).

B. Rice Transformation

RAmy1A/GUS plasmids were co-transformed into rice protoplasts (*Oryza sativa* L. japonica varieties, Nipponbare, Kinuhikari and Toride-1) by electroporation as previously described (Shimamoto, et al., 1989, Fertile transgenic rice plants regenerated from transformed protoplasts, *Nature*, 338:274-276 and Kyoizuka and Shimamoto, 1991 Transformation and regeneration of rice protoplasts, in *Plant tissue Culture Manual* (Lindsey, K. ed). Dordrecht:Kluwer Academic Publishers B1:1-16).

The *hph* gene (hygromycin phosphotransferase) was used as the selectable marker in these studies. Hygromycin B resistant calli were screened for GUS activity by incubating a portions of calli with X-glucuronide solution (Jefferson, 1987 *supra*). GUS positive calli were further cultured and plants were regenerated from these callus cultures.

C. Southern blot analysis

The GUS positive R1 plants, derived from two lines each of H4/GUS and E4/GUS primary transgenic lines, were used for Southern blot analysis. Total genomic DNA was isolated from mature leaves, digested by the restriction enzyme EcoRI and transferred onto a positively charged nylon membrane (Amersham). The coding region of the *gusA* gene was labeled and amplified with digoxigenin11-dUTP by polymerase chain reaction and used for probing the intact *RAMy1A/gusA* genes. Hybridization and chemiluminescence signal detection were performed according to manufacturer's specifications (Boeringer Mannheim Biochemica).

D. GUS assays

For histochemical analysis of GUS activity, germinating seeds were hand-cut with a razor and stained with X-glucuronide solution (5-bromo-4-chloro-3-indolyl glucuronide) as previously described in Kyojuka, et al., 1991 *supra* and Terada, et al., 1993, *Plant J.* 3:2412-252. For quantitative analysis, crude extracts from transgenic rice seeds were used for fluorometric assays of GUS activity as described previously (Kyojuka, et al., 1991; Terada, et al., 1993).

For developmental studies, transgenic R1 seeds were peeled off, sterilized with 10% NaOCl for 10 min and washed with distilled water. Seeds were germinated in plastic wells containing water for 2, 4, 6, and 8 days at 30°C under light.

For quantitative measurements of GUS activity germinating seeds were divided into the embryo and endosperm portions. In the case of the embryo, residual amount of endosperm, roots and shoots were removed before the assay.

For the analysis of hormonal regulation of the *RAmy1A/gusA* chimeric genes, transgenic R1 seeds were deembryonated and the embryoless seeds and sliced longitudinally into three pieces. Each slice was
5 treated with acetate buffer (10 mM sodium acetate pH 5.2), 10^{-7} M GA3 in acetate buffer, 10^{-7} M GA₃, and 10^{-5} M ABA in acetate buffer, for 4 days at 30C in the dark. Treatment slices were then used for the histochemical and the quantitative GUS assays.

10 E. RESULTS

1. *Southern blot analysis of transgenic rice plants*

Southern blot analysis confirmed the presence of
15 H4/GUS and E4/GUS gene fusions in the rice genome using the coding region of *gusA* as a probe (Figure 7A). The results indicated that complete sequences of the H4/GUS and E4/GUS chimeric genes were present in transgenic plants and that the *gusA* gene was
20 stably transmitted to the progeny (Figure 7B and 7C). In addition to the complete copies of the transgenes, several rearranged copies were also detected. The copy number of intact chimeric genes was estimated to be 1-3 per haploid genome.

25 2. *GUS activity in transgenic rice seeds*

To compare the relative expression levels of the H4/GUS and E4/GUS genes, the embryos and endosperms of germinated transgenic seeds were separated and the
30 GUS activity in each tissue was measured fluorometrically (Figure 8). GUS assays of endosperm tissue were performed at 6 days of germination, the time when α -amylase expression in the aleurone layer is at its highest (Figure 9).

Histochemical examination of 6-day germinated seeds showed that GUS activity was restricted to the scutellum of the embryo and the aleurone layer of the endosperm. In all four lines transformed with the H4/GUS gene, the aleurone activity was higher than the scutellum activity (Figure 8A). Similarly, four of the five lines transformed with E4/GUS gene showed GUS activity to be higher in the aleurone than in the scutellum (Figure 8B). Comparisons between H4/GUS and E4/GUS transformed lines, revealed no significant differences in GUS activity in the scutellar and aleurone tissues.

3. *Temporal and spatial regulation of RAMy1A/GUS expression during germination of transgenic rice seeds*

To investigate the role of the RAMy1A promoter in the temporal and spatial expression of heterologous genes during rice seed germination, histochemical and quantitative GUS assays on transgenic seeds were performed. Histochemical analysis of the H4/GUS chimeric gene showed that GUS activity could be detected in the scutellar epithelium after 2 days of germination and that this activity spread into the adjacent aleurone layer by day 4. On day 6, the GUS expression in the aleurone layer increased to the extent that it covered all portions of the seed.

To quantify the levels of GUS activity revealed by histochemical analysis, the GUS activities of germinating seeds derived from two H4/GUS transgenic lines (T21 and N33) were measured (Figure 9A). Scutellar GUS activity (open circles) appeared on day 2, peaked on day 4 and decreased thereafter.

In contrast, aleurone GUS activity (closed circles) was first detected on day 4, peaked on day 6

and decreased sharply by day 8. These results clearly show that the *RAmy1A/GUS* fusion genes are differentially regulated in scutellum and aleurone tissues during rice seed germination.

5 Similar experiments were performed on seeds from two lines of rice (K43 and T62) transformed with the E4/GUS gene fusion (Figure 1B). Histochemical assays revealed patterns of expression nearly identical to those observed for the H4/GUS gene. Quantitative
10 assays of GUS activity in the scutellar and the aleurone layers of germinated seeds indicated that the E4/GUS gene is expressed in the scutellum on day 2 and peaks at day 4. GUS activity in the aleurone layer is first detected at day 4 and peaks on day 6 (Figure 9B).
15 These results show that the -748 to +31 region in the H4/GUS gene and the -232 to +31 region in the E4/GUS gene, function identically with respect to the localization of expression and the developmental regulation during seed germination.

20

4. *Hormonal regulation of the RAmy1A/GUS genes in the aleurone layer of transgenic rice seeds*

25 To examine effects of exogenously GA and ABA on the expression of the H4/GUS and E4/GUS genes in seeds, deembryonated seeds were cut into three slices and each slice was treated with GA or a combination of GA and ABA. The histochemical examination of the H4/GUS seed slices treated with GA-free buffer showed
30 no GUS activity. However, seed slices treated with GA showed GUS activity in the aleurone layer. The observed induction of the GUS expression by GA was suppressed by addition of ABA. Similar GA induction of GUS in the aleurone layers of pE4/GUS transformed
35 seeds was also observed.

In an attempt to quantify GA induction of the *RAmy1A* promoter in transgenic seeds, GUS activity was measured for both GA, and GA+ABA treated seeds (Figure 10). A 10 to 40-fold increase in GUS activity was observed after GA treatment of H4/GUS (Figure 10A) and E4/GUS (Figure 10B) seeds. When ABA was added along with GA, the GUS activity was suppressed to a level just above background. The degree of GA induction and ABA suppression was similar for both H4/GUS and E4/GUS derived seeds.

While the invention has been described with reference to specific methods and embodiments, it will be appreciated that various modifications and changes may be made without departing from the invention.

78

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- 5 (i) APPLICANT:
- (A) NAME: The Regents of the University of
 California
- (B) ADDRESS: 300 Lakeside Drive
- (C) CITY: Oakland
- 10 (D) STATE: CA
- (E) COUNTRY: USA
- (F) POSTAL CODE: 94612-3550
- (ii) TITLE OF INVENTION: PROCESS FOR PROTEIN PRODUCTION
15 IN PLANTS
- (iii) NUMBER OF SEQUENCES: 7
- (iv) CORRESPONDENCE ADDRESS:
- 20 (A) ADDRESSEE: Dehlinger & Associates
- (B) STREET: 350 Cambridge Avenue, Suite 250
- (C) CITY: Palo Alto
- (D) STATE: CA
- (E) COUNTRY: USA
- 25 (F) ZIP: 94306
- (v) COMPUTER READABLE FORM:
- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- 30 (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: PatentIn Release #1.0, Version #1.25
- (vi) CURRENT APPLICATION DATA:
- (A) APPLICATION NUMBER:
- 35 (B) FILING DATE:
- (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
- (A) APPLICATION NUMBER: US 08/153,563
- 40 (B) FILING DATE: 16-NOV-1993

79

(viii) ATTORNEY/AGENT INFORMATION:

- (A) NAME: Fabian, Gary R.
(B) REGISTRATION NUMBER: 33,875
(C) REFERENCE/DOCKET NUMBER: 2000-0452.41

5

(ix) TELECOMMUNICATION INFORMATION:

- (A) TELEPHONE: (415) 324-0880
(B) TELEFAX: (415) 324-0960

10

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 820 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

15

(ii) MOLECULE TYPE: DNA (genomic)

20

(ix) FEATURE:

- (A) NAME/KEY: misc_feature
(B) LOCATION: 1..820
(D) OTHER INFORMATION: /standard_name= "HV-18"

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

30

GGATCCTAGC TACGGACAGC GCCCCGGTTA TGGAGGCCGA CAGCCGCGGC GCGCGGCTGC 60
GTAGCAGTGC AGCGTGAAGT CATAGATAGA CTGTAGAGGG CATGGCGGCA AGTGAAAACA 120
CACTTCGGTT TGTTCGTGTG AGTCAGTTGG ATCTGCTTTG GCCTGGCGAT AACGTCTCCG 180
GCCATTGTTT ATCACGGCGC CTGCTTATCC CTCGAAAGT TTGAGCAAAA GGTGCAGCTT 240
CTTCTAGTA CAGAAATGAC GTCCAGAGTT GCAGCAACCC ATTCGGAACCT CCTGGTGGAT 300
GCCAACGAAA TTAAATGGGA TAAACTTAG TGAAGAATCT ATATTTTCTT GCAACAACAT 360
ACTCCTACCC TCACGAATTG AATGCTCATC GAACGAATGA ATATTGAT ATATGTTGAT 420

40

80

CTCTTCGGAC TGAAAAAGTT TGAAC TCGCT AGCCACAGCA CACTATTCCA TGAAAAATGC 480
TCGAATGTTT TGTCTAGAA AAACAGAGGT TGAGGATAAC TGACGGTCGT ATTGACCGGT 540
5 GCCTTCTTAT GGAAGGCGAA GGCTGCCTCC ATCTACATCA CTTGGGCATT GAATCGCCTT 600
TTGAGCTCAC CGTACCGGCC GATAACAAAC TCCGGCCGAC ATATCCACTG GCCCAAAGGA 660
GCATTCAAGC CGAGCACACG AGAAAGTGAT TTGCAAGTTG CACACCGGCA GCAATTCCGG 720
10 CATGCTGCAG CACACTATAA ATACTGGCCA GACACACAAG CTGAATGCAT CAGTTCTCCA 780
TCGTACTCTT CGAGAGCACA GCAAGAGAGA GCTGAAGAAC 820

15 (2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2389 base pairs
(B) TYPE: nucleic acid
20 (C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

25

(ix) FEATURE:

- (A) NAME/KEY: misc_feature
(B) LOCATION: 1..2389
(D) OTHER INFORMATION: /standard_name= "RAMY-1A"
30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

CTGCAGGCAT GCGAGAGGCA CGGGGTTTGA TCCCCCGCT CTCCATCGGC ACTGTTTTTT 60
35 AACATCAAAC GCTGTTGAT CCACTATCTG TTAATTTGCG AAACACAAC AAATCTTTTT 120
TTTTTTTTGC CGGTGCGTGC AGTGTGACGT CCAAGGCATG GCGCATGGC GCCTCCCCTC 180
40 TTTCCCTTGA TCTTTTCATC AGTTGTTCT TCTTGCAGAA AAGCTGTTCT GTTAAGTCGG 240

81

TTCCGATCTG CTCTTGGGCT CTTGCCAGAA ACAACCTGTG TACGCCAGAC TTATCAAGCC 300
AACCATCCTG ATGAGCCTCT GCTTATACAA GCCTTTGACT CCAAAAAGGA CGAGGAGGCT 360
5 TGCAGCCGCA CGGAAATAAG CCGACCGATC CTTTATTGCT CTATCTTTTT CCCTTGGAAT 420
AAAAACAGC CCAATTAAAA TCTGGGATGA AACTATGGCT AGCTGTTCCG GGTGTCAGTT 480
CTCGGGACGC TACCGTTGTT TTGTTTGAAC CGGAATGTTT AGGGCGGTTC ACACCATAGA 540
10 CTTGGAGCCA AGTGGTTCCA TCCACAAAAT TTTCTCATCT TGAATATTCT GTTATCTGCC 600
TCGACAGACG CACCATATCC TGTGTTCCAG AATGAATGTG CTACAGCCAA CGTGCTGCAT 660
15 GAAATTGCT GAAATCGTGC TAAAATGTGC ATGGCAACAG GAACCTGATG CCCTGGTCCT 720
GTGGAAGTGC CACGGGAAAG TATTTTTTAT AGCTAGGTGC AATCGTATCT AGGTGTATAC 780
ATGTCACCTA CATAGCTACT CCCCTTTATC TTAAATATA ATAATTTTAA ACTCTCAGTA 840
20 TTTGTCCTAA AATATAACAA ATTCTCCATC AACATTATCT TCCCAACCAA TCACAACCCT 900
TCATCATTAA TTTTTTCCCC CTACCTCCAC TACTCATCTA ATCACAACCC TCCAACACTC 960
25 ACTTCTATCT ACTTTCTTAA TAACTGTCTT CAACCCTAAA ACTTCTTATA TTTTAGGACG 1020
GAGGGAGTAT CTAAATATTT CATAAAAAAA ATGTTAAGAT AGATAAAGAA GATATAAACC 1080
CACTATGCAA ACATGCACAT CAAAATTTAA TTTACAGTAA AGAAACAGAA ATAACATATT 1140
30 CTATTTGTGC TGGAGATGTA CTGTTACAA TATTGTTTTT TTATTTTTTA TTTATCTGAT 1200
TATATATCTG TTTCAGCCTT GCATGGTTGT GTATGTTTGT GTATAGACTT ATGCCATTGT 1260
35 GATTGATGCT ACCAATTATT TTCAGACTAT TTTTTTATAG AGGAATTTTA TAGTTCTTGA 1320
GAAAATACCT TGAAGTATCT AAATTTTACA CTAAAATTGT TGGTACCTTG AGGTACAAAG 1380
TACCTAGAGG TACCAAATTT TACTAGAAAA TTGTGGCACC TTAGGTACC TTCTCAAAAA 1440
40 TAGTACAATT ATGGGCCGTT TTGGATTTAG TGCCAAAACG TGCTCTACAA ATATTTTGAT 1500

82

AGTTTGAACA GTGCATAAGA CGGGTTTGGT TTGAAGCCAA ATCATTGGCA TTGCCAATGT 1560
CCAATTTGAT ATTTTCTATA TTATGCTAAA AGCTTGGTTC TAAATTGGCC TCCAACCAAA 1620
5 TACAACTCTA CTCTACCAAA AAATTTGTAG TGCCAAAAC TGCCTAGGTT TTGTCACTAC 1680
CAACATTTTG GTAAGTATTA AACCAAACAA GCCCTACATT TTTTATGTA CATTTAAGTT 1740
GTATGTAAAT GATGGGTGCG GTTGACCTA GGTGAAAAA AATACATATT CGCCACAAC 1800
10 CGCAACATGT ACCAATTCAG CAGCAAGTGT AAGAGAGAAG ATTTCTCTCG TTTTACACGC 1860
GCACGTTCAA TTCCTGAACT ACTAAACGGT ATGATTTTTT GCAAAAATT TCTATAGGAA 1920
15 AGTTACTTAA AAATTATATT AATCTATTTT TAAAATTTAA AATAGTTAAT ACTCAATTAA 1980
TTATACGTTA ATGGCTCAGC TCGTTTTGCG TACATTCTCA ATCGATTCTT TTCCTCTGCT 2040
CTCAAATGCT CTGTGTGCGA TCAGGTATTC ATGTTGAGCT CGCACAAGCA CAAGCAAGAC 2100
20 AGATGGAATT CCTACTGACC TCGGCCTTTT GAGTCGCTCC AACTCTCAA GTCTCAAGGC 2160
CATTAAATTG CCTATGGGCT CACCAGCCAA TAACAAACTC CGGCTGTTAT CCATCCAATC 2220
25 CAGTGTCCCA AAGCAACATT CAAGCCCAGC CAGGCCTCCA AAAGTTGCAA GTTGAGCATG 2280
GCAAAATCCC CGGCAATTCT CGACTATAAA TACCTGACCA GACACACCCA GGAGCTTCAT 2340
CAATCATCCA TCTCCGAAGT GTGTCTGCAG CATGCAGGTG CTGAACACC 2389
30

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- 35 (A) LENGTH: 31 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

40

83

(ix) FEATURE:

(A) NAME/KEY: misc_feature

(B) LOCATION: 1..31

(D) OTHER INFORMATION: /standard_name= "31 bp RAmy3D"

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

GAGACCGGGC CCGACGCGG CCGACGCGGC G

31

10

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 31 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

15

(ii) MOLECULE TYPE: DNA (genomic)

20

(ix) FEATURE:

(A) NAME/KEY: misc_feature

(B) LOCATION: 1..31

(D) OTHER INFORMATION: /standard_name= "31 bp RAmy3E"

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

GAGAGCTCGC GCCGCCTCGA TCGGCGCGGC G

31

30

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 11 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

35

(ii) MOLECULE TYPE: DNA (genomic)

40

84

(ix) FEATURE:

(A) NAME/KEY: misc_feature

(B) LOCATION: 1..11

(D) OTHER INFORMATION: /standard_name= "11 bp RAmy3D"

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

TTCCGGCTTG C

11

10

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 11 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

15

(ii) MOLECULE TYPE: DNA (genomic)

20

(ix) FEATURE:

(A) NAME/KEY: misc_feature

(B) LOCATION: 1..11

(D) OTHER INFORMATION: /standard_name= "11 bp RAmy3E"

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

TTGCGGCTTG C

11

30

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 12 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

35

(ii) MOLECULE TYPE: DNA (genomic)

40

85

(ix) FEATURE:

(A) NAME/KEY: misc_feature

(B) LOCATION: 1..12

(D) OTHER INFORMATION: /standard_name= "Taka-amylase"

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

CGGCCCGTCG GC

12

IT IS CLAIMED:

1. A method of producing a protein in a monocot
plant seed, where the seed contains an endosperm
5 surrounded by an aleurone or scutellar epithelium layer,
comprising
malting seeds containing a chimeric gene having (i)
a transcriptional regulatory region inducible during seed
germination, (ii) a heterologous first DNA sequence
10 encoding said protein, and (iii) a second DNA sequence
encoding signal polypeptide where said second DNA
sequence is operably linked to said transcriptional
regulatory region and said first DNA sequence, and where
said signal polypeptide is in translation-frame with said
15 protein and is effective to facilitate secretion of said
protein across said aleurone or scutellar epithelium
layer into the endosperm,
where said seeds are malted under conditions to
induce expression of said transcriptional regulatory
20 region and production of said protein.
2. A method of claim 1, wherein the seed is from a
cereal plant.
- 25 3. A method of claim 2, wherein the cereal seed is
selected from the group consisting of wheat, rice, oats,
rye, corn, sorghum, millet and barley.
4. A method of claim 1, wherein the cereal seed is
30 rice.
5. A method of claim 1, wherein the
transcriptional regulatory region includes a promoter
region selected from the group of genes consisting of an

α -amylase gene, a sucrose synthase gene and a sucrose-6-phosphate synthetase gene.

5 6. A method of claim 5, wherein the transcriptional regulatory region is obtained from an α -amylase gene.

10 7. A method of claim 6, wherein the α -amylase gene is selected from the group consisting of RAMy1A (SEQ ID NO:2), RAMy3B, RAMy3C, RAMy3D, HV18 (SEQ ID NO:1) and RAMy3E.

15 8. A method of claim 6, where said promoter is from the RAMY3D gene or homologs thereof.

9. A method of claim 1, where expression mediated by said transcription regulatory region is specifically regulatable by a small molecule.

20 10. A method of claim 9, where said small molecule is a plant hormone or cytokine.

25 11. A method of claim 10, where said small molecule is selected from the group consisting of abscisic acid, gibberellic acids, indoleacetic acids, kinetins, butyric acid, oxalic acid, acetic acid, okadaic acid and arachidonic acid.

30 12. A method of claim 10, where said transcription regulatory region is from the RAMY3D gene, or a homolog thereof, and said small molecule is abscisic acid.

13. A method of claim 9, wherein the malting step further comprises the addition or removal of said small

molecule in an amount effective to stimulate production of the protein.

5 14. A method of claim 9, wherein the transcriptional regulatory region is obtained from a sucrose synthase gene or sucrose-6-phosphate synthetase gene.

10 15. A method of claim 14, wherein said small molecule is a sugar or sugar-phosphate derivative.

 16. A method of claim 15, where said sugar is glucose or sucrose.

15 17. A method of claim 1, wherein the protein is selected from the group consisting of enzyme, antibody, growth factor, cytokine, hormone, and antigen.

20 18. A method of claim 1, where said protein is selected from the group consisting of α -antitrypsin, antithrombin 3, fibrinogen, human serum albumin, factor VIII, granulocyte colony-stimulating factor and granulocyte macrophage colony-stimulating factor.

25 19. A method of claim 1, where said protein is an industrial protein.

30 20. A method of claim 19, where said industrial enzyme is selected from the group consisting of xylanase, oxidoreductase, peroxidase, glucanase, α -amylase, phytase and glucose oxidase.

 21. A method of claim 1, wherein the signal polypeptide is from a cereal gene selected from the group

consisting of an α -amylase gene, a sucrose synthase gene and a sucrose-6-phosphate synthetase.

5 22. A method of claim 21, wherein the signal polypeptide is from an α -amylase gene.

23. A method of claim 22, wherein the amylase gene is RAMY1A, RAMY3B, RAMY3C, RAMY3D, HV18 or RAMY3E.

10 24. A method of claim 1, where following said malting, said protein is isolated in a substantially pure form.

15 25. A method of claim 1, where said seed contains an embryo, and wherein the embryo is (i) separated from the other seed parts, and (ii) germinated under conditions that induce expression of said transcriptional regulatory region and production of said protein.

20 26. A method of claim 25, where said seeds contain a second chimeric gene having (i) a transcriptional regulatory region inducible during seed germination, (ii) a heterologous DNA sequence that encodes a polypeptide, where said DNA sequence is operably linked to said
25 promoter, and

said separated embryos are germinated under conditions that (i) induce expression of the transcriptional regulatory region of the second chimeric gene, and (ii) production of said polypeptide of the
30 second chimeric gene.

27. A method of claim 26, where the transcriptional regulatory region of the second chimeric gene is obtained from a gene selected from the group of genes consisting

of an α -amylase gene, a sucrose synthase gene and a sucrose-6-phosphate synthetase gene.

28. A method of claim 1, where said seed contains
5 an embryo and where said malting includes,
separating embryos of the seed from the endosperm;
contacting the endosperm with the plant hormone
which induces expression of the expression cassette; and
isolating the heterologous protein from the
10 endosperm.

29. A mash product produced from the malted seeds
of claim 1.

15 30. A mash product of claim 29, where said
polypeptide is useful for veterinary applications.

31. A malted seed product produced from the malted
seeds of claim 1.
20

32. A malted seed product of claim 31, where said
polypeptide is useful for veterinary applications.

33. A germinated seed produced by malting the seeds
as in claim 1.
25

34. A chimeric gene of any of claims 1, 5 through
10, and 17 through 23.

35 35. A plant expression vector comprising the
chimeric gene of claim 34.

36. A plant cell comprising the chimeric gene of
claim 34.
35

37. A plant cell of claim 36, where said plant is selected from the group consisting of rice, wheat, oats, rye, corn, sorghum, millet, and barley.

5 38. A transgenic plant comprising the plant cell of claim 36.

 39. Seeds produced by growing the transgenic plant of claim 38.

10

 40. Seeds of claim 39, where said transgenic plant is selected from the group consisting of rice, wheat, oats, rye, corn, sorghum, millet, and barley.

15 41. A chimeric gene, comprising
 (i) a transcription regulatory region inducible during seed germination, where expression mediated by said region is specifically regulatable by a small molecule, and (ii) a heterologous DNA sequence that
20 encodes the polypeptide, where said DNA sequence is operably linked to said promoter.

 42. A chimeric gene of claim 41, wherein the transcriptional regulatory region includes a promoter
25 region selected from the group of genes consisting of an α -amylase gene, a sucrose synthase gene and a sucrose-6-phosphate synthetase gene.

 43. A chimeric gene of claim 42, wherein the transcriptional regulatory region is a promoter of an α -
30 amylase gene.

 44. A chimeric gene of claim 43, wherein the α -amylase gene is selected from the group consisting of

RAMy1A (SEQ ID NO:2), RAMy3B, RAMy3D, HV18 (SEQ ID NO:1) or RAMy3E.

5 45. A chimeric gene of claim 44, where said promoter is from the RAMY3D gene or homologs thereof.

10 46. A chimeric gene of claim 41, where said DNA sequence encodes a polypeptide selected from the group consisting of enzyme, antibody, growth factor, hormone, cytokine, and antigen.

15 47. A chimeric gene of claim 41, where said polypeptide is selected from the group consisting of α -antitrypsin, antithrombin 3, fibrinogen, human serum albumin, factor VIII, granulocyte colony-stimulating factor and granulocyte macrophage colony-stimulating factor.

20 48. A chimeric gene of claim 41, where said polypeptide is an industrial protein.

25 49. A chimeric gene of claim 48, where said industrial enzyme is selected from the group consisting of xylanase, oxidoreductase, peroxidase, glucanase, α -amylase, phytase and glucose oxidase.

30 50. A chimeric gene of claim 41, wherein the transcriptional regulatory region is a promoter of a sucrose synthase or a sucrose-6-phosphate synthetase gene.

35 51. A chimeric gene of claim 41, where said chimeric gene further encodes a signal sequence, where a second DNA sequence encoding the signal sequence is operably linked to said transcription regulatory region

and said heterologous DNA sequence, and where said signal sequence is (i) in translation-frame with said polypeptide encoded by the heterologous DNA sequence, and (ii) effective to facilitate secretion of said polypeptide from the cell.

52. A chimeric gene of claim 51, wherein the signal sequence is from a cereal gene selected from the group consisting of an α -amylase gene, a sucrose synthase gene and a sucrose-6-phosphate synthetase.

53. A chimeric gene of claim 52, wherein the signal sequence is from an α -amylase gene.

54. A chimeric gene of claim 53, wherein the α -amylase gene is selected from the group consisting of RAmy1A, HV18, RAmy3D and RAmy3E.

55. A plant expression vector comprising the chimeric gene of any of claims 41-54.

56. A plant cell comprising the chimeric gene of any of claims 41-54.

57. A plant cell of claim 56, where said plant is selected from the group consisting of rice, wheat, oats, rye, corn, sorghum, millet, and barley.

58. A transgenic plant comprising the plant cell of claim 56.

59. A transgenic plant of claim 58, where said plant is selected from the group consisting of rice, wheat, oats, rye, corn, sorghum, millet, and barley.

60. A method for modulating expression of a polypeptide in plant tissue cell culture, where said cells are from a monocot plant, comprising
5 culturing the plant cells of claim 56 under conditions that facilitate plant cell growth, and modulating expression mediated by the transcription regulatory region by addition or removal of said small molecule to the plant cell culture.

10 61. A method of claim 60, wherein the cells are derived from scutellar epithelium of cereal plants.

62. A method of claim 61, wherein the cereal is selected from the group comprising wheat, rice, oats,
15 rye, corn, sorghum, millet and barley.

63. A method of claim 62, wherein the cereal is rice.

20 64. A method of claim 60, where said modulating is an inhibition of expression, and said small molecule is a sugar or sugar-phosphate derivative.

25 65. A method of claim 64, where said sugar is glucose or sucrose.

66. A method of claim 60, where said modulating is an induction of expression, and said small molecule is a plant hormone.

30 67. A method of claim 66, where said small molecule is selected from the group consisting of abscisic acid, gibberellic acids, indoleacetic acids, kinetins, butyric acid, oxalic acid, acetic acid, okadonic acid and
35 arachidonic acid.

68. A method of claim 60, wherein the culturing step further comprises the addition or removal of sugar or sugar-derivatives in an amount effective to stimulate production of the heterologous protein.

5

69. A method of claim 68, wherein the transcriptional regulatory region is a promoter of a sucrose synthase or sucrose-6-phosphate synthetase gene.

10

70. A method of claim 60, where said modulation is the enhancement of expression of the polypeptide in plant tissue cell culture, said enhancement is accomplished by addition of said small molecule to the plant cell culture, where said small molecule specifically induces expression from the transcription regulatory region.

15

71. A method of claim 70, where said small molecule is a plant hormone or cytokine.

20

72. A method of claim 71, where said transcription regulatory region is from the RAMY3D gene, or a homolog thereof, and said small molecule is abscisic acid.

25

73. A method for producing a polypeptide in plants cells, comprising culturing the plant cell of claim 56 *in vitro* under conditions that facilitate plant cell growth and supplying a small molecule to the cells that specifically induces production of said polypeptide.

30

74. A method of claim 73, where said producing further includes isolating said polypeptide in a substantially pure form.

35

75. A method of producing a polypeptide in plant cells, comprising

culturing the plant cell of claim 56 *in vitro*
(i) under conditions that facilitate plant cell growth,
and (ii) in the presence of a small molecule that
inhibits expression from the transcription regulatory
5 region, where said small molecule is present at a
concentration effective to repress expression from the
transcription regulatory region, and

reducing the concentration of said small
molecule in the culture to a level that permits
10 expression from the transcription regulatory region and
allows production of said polypeptide.

76. A method of claim 75, where said small molecule
is a sugar or sugar-phosphate derivative.
15

77. A method of claim 76, where said sugar is
glucose or sucrose.

78. A method of claim 75, where said transcription
20 regulatory region is the promoter of the RAMY3D gene.

79. A method of claim 78, where after said
reducing, said culture further includes addition of a
second small molecule that induces expression of the
25 RAMY3D promoter.

80. A method of claim 79, where said second small
molecule is abscisic acid.

81. An isolated DNA sequence essentially consisting
30 of the nucleic acid sequence of SEQ ID NO:1 or SEQ ID
NO:2.

82. An isolated DNA sequence essentially consisting of a transcriptional regulatory region obtained from SEQ ID NO:1 or SEQ ID NO:2.

5 83. An isolated DNA sequence of claim 82 operably linked to a heterologous DNA sequence encoding a protein.

10 84. An isolated DNA sequence of claim 83 that further includes a second DNA sequence encoding signal polypeptide where said second DNA sequence is operably linked to said transcriptional regulatory region and said heterologous DNA sequence, and where said encoded signal polypeptide is in translation-frame with said encoded protein.

1/11

FIG. 1A



RAmy3D PROMOTER/GUS FUSION

FIG. 1B

RAmy1A PROMOTER/GUS FUSION

2/11

FIG. 2





FIG. 4



3/11

FIG. 3B





FIG. 3A



4/11

FIG. 5




FIG. 6B

RAmy 3D	-264	GAGAC <u>CGGG</u> CCCCGACGCGGCCGACGCGGGCG (SEQ ID NO: 3)	
		++++ + ++ +++ + ++ ++ ++++++	
RAmy 3E	-392	GAGAGCTCGCGCCGCCTCGATCGGCGCGGGCG (SEQ ID NO: 4)	
RAmy 3D	-648	TTCCGGCTGC	(SEQ ID NO: 5)
		++ ++++++	
RAmy 3E	-479	TTGCGGCTTGC	(SEQ ID NO: 6)

5/11






FIG. 6A(1)

FIG. 6A(2)

FIG. 6A(3)

6/11



7/11

FIG. 8B





FIG. 8A



8/11

FIG. 9A(2)





FIG. 9A(1)



9/11

FIG. 9B(2)





FIG. 9B(1)



10/11

FIG. 10A(2)





FIG. 10A(1)



11/11

FIG. 10B(2)




FIG. 10B(1)

